3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643 (310) 370-8370 (310) 370-7026 FAX www.hygienetech.com

August 8, 2008

California State Board of Equalization 450 N Street Sacramento, California 94279

Document No. 20803001.6

Attention:

David Gau

Regarding:

Limited Indoor Air Quality Survey

2ND Floor

Dear Mr. Gau:

On various dates in March and April of 2008, industrial hygienists with Hygiene Technologies International, Inc. (HygieneTech) conducted a limited indoor air quality survey on the 2ND Floor of the California State Board of Equalization building located at the above mentioned address. At the time of the survey, various samples were collected and direct-reading instruments were used to assess the general indoor air quality, with a clear emphasis on establishing fungal growth exposure potential data. I have enclosed our report, which included general observations, sample and direct-reading results, a discussion of the data, conclusions, and recommendations.

If you have any comments or questions regarding the information contained in this report, please do not hesitate to contact our offices directly at (310) 370-8370.

Sincerely,

HYGIENE TECHNOLOGIES INTERNATIONAL, INC.

Brian P. Daly, CIH, PE

President

3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643 (310) 370-8370 (310) 370-7026 FAX www.hygienetech.com

LIMITED INDOOR AIR QUALITY SURVEY

450 N STREET – 2ND FLOOR SACRAMENTO, CALIFORNIA

PREPARED FOR:

CALIFORNIA STATE BOARD OF EQUALIZATION
450 N STREET
SACRAMENTO, CALIFORNIA

PREPARED BY:

HYGIENE TECHNOLOGIES INTERNATIONAL, INC. 3625 DEL AMO BOULEVARD, SUITE 180 TORRANCE, CALIFORNIA

AUGUST 8, 2008



1.0 BACKGROUND

On various dates in March and April of 2008, industrial hygienists with Hygiene Technologies International, Inc. (HygieneTech) conducted a limited indoor air quality survey on the 2^{ND} Floor of the California State Board of Equalization Building located at 450 N Street in Sacramento, California. During the survey, a variety of samples were collected and direct-reading instruments were used to assess the general indoor air quality on the 2^{nd} Floor of the subject building. Various air and surface samples were collected in order to assess fungal growth exposure potentials and to establish fungal growth assessment information on selected building material surfaces. In addition, air samples were collected throughout the floor for fibrous dust, microbial volatile organic compounds (MVOCs), and total dust analysis and direct-reading instruments were used to determine airborne volatile organic compounds (VOCs), carbon dioxide (CO₂), ozone (O₃), air temperature, and relative humidity.

2.0 OBSERVATIONS

The interior building materials of 2ND Floor included, but were not limited to, metal window frames; painted gypsum board and/or metal window sills; metal doorjambs and door frames; painted gypsum board walls in the general work areas; tile covered walls and painted gypsum board ceilings in the restrooms; suspended 2' by 4' ceiling tiles in the general work areas; vinyl cove base; carpet flooring in the general work areas; and ceramic or vinyl tile flooring in the restrooms and break rooms.

The furnishings in the surveyed areas included desks, upholstered chairs, shelves, fabric covered cubicles, office supplies, computers, and other electronic office equipment. The furnishings did not appear to support fungal growth, nor did they appear to have been affected in any other manner by water intrusion. However, be advised that visible accumulation of debris, dust, and other particulates was observed on the reverse side of all sampled HVAC supply air registers.

3.0 SAMPLING AND ANALYSIS

Air samples were collected and subsequently analyzed for fungi (including yeasts, molds, rusts, smuts, and mushrooms) by trained and experienced microbiologists at laboratories accredited by the American Industrial Hygiene Association (AIHA) and that successfully participate in the AIHA Environmental Microbiology Proficiency Analytical Testing (EMPAT) Program. Other samples were collected for airborne fibers, MVOCs, and total dust determinations using SKC® brand Airchek® 52 sampling pumps and the appropriate sampling media. Pump flow rates were established and verified using a BIOS DryCal DC-Lite primary flow meter. Those samples were collected and analyzed along with blanks (identical sampling media through which no air was drawn), when necessary, at laboratories accredited by the American Industrial Hygiene Association (AIHA) through successful participation in the National Institute for Occupational Safety and Health (NIOSH) Proficiency Analytical Testing Program. Direct-reading instruments were used to determine airborne O₃ and VOC levels, the results of which appear in Table 20803001-144 in Appendix A of this report. A discussion of the airborne CO₂ data, along with air temperature and relative humidity results, appears in Section 4.0 of this report. Additional information concerning the specific sampling and analytical methods appears below.



3.0 SAMPLING AND ANALYSIS (CONTINUED)

3.1 Airborne Total Fungi

Air samples for airborne total (viable and nonviable) fungi determinations were collected using a Zefon brand Bio-Pump[™] equipped with Allergenco-D[™] cassettes. All such samples were collected at various indoor locations and two samples were collected outdoors on the applicable survey dates for comparison purposes. The resultant data, which are presented in spores per cubic meter of air (spores/M³), appear in Table 20803001-138.

3.2 <u>Airborne Viable Fungi</u>

Air samples for airborne viable fungi determinations were collected on malt extract agar (MEA) using a Gast brand high volume air-sampling pump equipped with an Aerotech 6[™] Single Stage Bioaerosol Sampler. Two outdoor samples were also collected on the applicable survey date for comparison purposes. The media was incubated prior to enumeration of colony-forming units per agar plate and the resultant data, presented in colony forming units per cubic meter of air (CFU/ M³), can be found in Table 20803001-139.

3.3 Surface Fungal Growth Potentials

Surface samples were collected for fungal growth assessment using Scotch® brand cellophane tape segments affixed to microscope slides. Additionally, surface fungi samples were collected from various heating, ventilating, and air conditioning (HVAC) supply air register surfaces using Healthlink® Transporters™ (Rayon tipped swabs immersed in 0.5 ml modified Stuart's transport medium). These data are presented in Table 20803001-140.

3.4 Airborne Fibrous Dust

Area air samples for fibrous dust were collected at stationary locations on 25-millimeter diameter, 0.8-micrometer pore size, mixed cellulose ester filters. The samples were analyzed by phase contrast microscopy (PCM) in accordance with the NIOSH Method 7400. These data are presented in fibers per cubic centimeter (f/cc) of air in Table 20803001-141.

3.5 Airborne Total Dust

Area air samples for total dust determination were collected at stationary locations on filter cassettes containing pre-weighed 37-millimeter diameter, polyvinyl chloride filters having a pore size of five micrometers. The samples were analyzed by gravimetric method in accordance with the NIOSH Method 0500. These data are presented in milligrams per cubic meter of air (mg/M³) and appear in Table 20803001-142.

3.6 <u>Microbial Volatile Organic Compounds</u>

Area samples for MVOCs were collected on solid sorbent tubes equipped with Sagelock fittings. The samples were analyzed by gas chromatography/ mass spectrometry, modified for MVOCs following the AIHA field guide. These data are presented in mg/M³ and appear in Table 20803001-143.



3.0 SAMPLING AND ANALYSIS (CONTINUED)

3.7 Airborne Volatile Organic Compounds

Direct-reading air measurements for VOCs were also recorded at various locations on the 2nd Floor using a RAE Systems, Inc. Mini-RAE 2000 photoionization detector, which is capable of detecting a wide variety of unsaturated hydrocarbons at airborne concentrations ranging from 0.1 to 10,000 parts per million (ppm). Prior to the survey, this instrument was calibrated using a 100-ppm isobutylene gas standard. These data are presented in parts per million (ppm).

3.8 <u>Airborne Ozone</u>

Direct-reading air measurements for O_3 were recorded at various locations using a Dräger colorimetric detector tube apparatus with the appropriate detector tubes. The data are presented in ppm.

3.9 <u>Airborne Carbon Dioxide</u>

Direct-reading air measurements for airborne CO₂ concentration was recorded at a stationary location using a Telaire[®] 7001 Carbon Dioxide and Temperature Monitor along with HOBO[®] data logger. The data are presented in ppm.

3.10 Air Temperature and Relative Humidity

Air temperature and relative humidity data were recorded at a stationary location using a Telaire® 7001 Carbon Dioxide and Temperature Monitor along with the HOBO® data logger.

4.0 DISCUSSION

4.1 <u>Airborne Total Fungi</u>

The airborne total fungi data showed common fungal spore types outdoors such as *Alternaria*, *Arthrinium*, ascospores, basidiospores, *Bipolaris/Drechslera* group, *Cladosporium*, colorless spores typical of *Penicillium* and *Aspergillus* species, *Curvularia*, *Epicoccum*, *Nigrospora*, *Oidium*, rusts, smuts, *Torula*, and/or unidentified mitosporic fungi, with *Cladosporium* predominating in all the samples. Additionally, less common fungal spores of *Chaetomium* and *Stachybotrys* were also found outdoors. Indoors, the ambient data showed low airborne concentrations of common fungal spores that included one or more of the following: *Alternaria*, ascospores, basidiospores, *Chaetomium*, *Cladosporium*, colorless spores typical of *Penicillium* and *Aspergillus* species, *Oidium*, rusts, smuts, *Torula*, and/or unidentified mitosporic fungi. The overall spore counts within the tested areas were well below the overall data recorded outdoors.

Additionally, the air samples collected within the ceiling plenums indicated generally low levels of fungal spore types such as *Alternaria*, *ascospores*, basidiospores, *Botrytis*, *Cladosporium*, colorless spores typical of *Penicillium* and *Aspergillus* species, *Oidium*, rusts, smuts, *Stachybotrys*, *Stemphylium*, *Torula*, *Ulocladium*, and/or unidentified mitosporic fungi, with the exception of two somewhat higher levels of *Cladosporium* recorded near Cubicles 37 and 55. In addition, the low



4.0 DISCUSSION (CONTINUED)

4.1 Airborne Total Fungi (Continued)

but detectable level of *Stachybotrys* recorded in one sample was likely an anomaly and therefore considered unremarkable. However, note that the airborne fungi results for the occupied space would suggest that such conditions in the ceiling plenums did not appear to have adversely affected the indoor air quality on the 2ND Floor. These data are not believed to pose a health risk beyond that posed by the outdoor environment where exposures to airborne fungi are expected.

4.2 Airborne Viable Fungi

The viable fungi data recorded outdoors showed overall levels of 1,239 CFU/M³ and 1,593 CFU/M³ in the two samples collected, with *Cladosporium* predominating in both samples. Indoors, low levels of common fungi were found, including *Alternaria*, *Aspergillus fumigatus*, Arthrosporeformer, *Cladosporium*, non-sporulating fungi, *Penicillium*, and/or yeasts. Again, the recorded data were unremarkable and are not believed to pose a health risk beyond that posed by the outdoor environment where exposures to airborne fungi are expected.

4.3 Surface Fungal Growth Potentials

The surface assessment data involving the samples collected from various cubicle partitions and other surfaces throughout the 2ND Floor indicated no evidence of fungal growth or above-background levels of loose fungal spores on those surfaces. Additionally, the surface assessment data from the samples collected from the HVAC supply air registers throughout the floor indicated loose fungal spores involving *Alternaria*, *Cladosporium*, *Penicillium/Aspergillus* types, and/or *Ulocladium* on six of the eight registers sampled. Be advised that visible accumulation of debris, dust, and other particulates was observed on the reverse side of all sampled HVAC supply air registers, and that such conditions are indicative of an environment that may promote fungal growth.

4.4 Airborne Fibrous Dust

The data recorded in the surveyed areas indicated that airborne fibrous dusts were detected at levels ranging from 0.007 to 0.012 f/cc. Because the samples were collected at stationary locations at approximate breathing zone height, the resultant data are expected to represent building occupant *exposure potentials* for those persons working in or passing through the areas monitored. These data, which are expected to represent employee *exposure potentials* to fibers of various types, including man-made and natural mineral fibers, cellulosics (paper or wood composition), gypsum, and other fibrous dusts common in the environment, are well below the current Cal-OSHA 8-hour TWA PEL for asbestos fibers of 0.1 f/cc, the most restrictive exposure limit for fibrous dusts.

4.5 <u>Airborne Total Dust</u>

Common dust that is typically identified in buildings usually contains a wide variety of materials including, but not limited to, gypsum crystals, cellulosic particles, fiberglass fragments, mineral grains from soil, fungi spores, fine glass fibers, textile and wood fibers, iron or steel fragments, dead skin cells, insect parts, animal dander, and pollens. Generally, exposure to low levels of such materials does not produce ill effects in most persons. In fact, these so-called *nuisance dusts* have a long



4.0 DISCUSSION (CONTINUED)

4.5 <u>Airborne Total Dust</u> (Continued)

history of little adverse effect to the lungs and are not known to produce significant diseases or toxic effects, such as collagen (scar tissue) formation, when exposure are kept under reasonable control.

The data recorded in the surveyed areas showed that airborne total dust was not detected above the laboratory detection limit of 0.17 mg/M³. Because the samples were collected at stationary locations at approximate breathing zone height, the resultant data are expected to represent building occupant *exposure potentials* for those persons working in or passing through the areas monitored. These data are well below the State of California, Department of Industrial Relations, Division of Occupational Safety and Health (Cal-OSHA) 8-hour time-weighted average (TWA) permissible exposure limit (PEL) for total dust of 10 mg/M³, as defined in Title 8 of the California Code of Regulations, Section 5155 (T8, CCR § 5155). Note that these data are also well below the American Conference of Governmental Industrial Hygienists 8-hour TWA threshold limit value (TLV-TWA) for particulate (not otherwise classified) of 10 mg/M³; the U.S. Environmental Protection Agency (EPA) National Ambient Air Quality Primary Standard of 0.26 mg/M³ (24-hour standard); and the American Society of Heating, Refrigerating, and Air-Conditioning Engineers, Inc. (ASHRAE) theoretical value for non-occupational environments of 1/10 of the TLV.

4.6 <u>Airborne Microbial Volatile Organic Compounds</u>

Microbial Volatile Organic Compounds (MVOCs) are composed of low molecular weight alcohols, aldehydes, amines, ketones, terpenes, aromatic and chlorinated hydrocarbons, and sulfur-based compounds that are known to be byproducts of microbial metabolism. MVOCs have a very low odor threshold, thus, making them easily detectable by smell. They often have strong odors and are responsible for the smells generally associated with fungal growth.

The airborne MVOC data indicated the presence of 1-butanol at levels ranging from 525 ng/M³ to 548 ng/M³. Microbial growth related VOCs would not be expected to be present indoors without additional MVOCs such as ethanol, 1-octen-3-ol, 2-octen-1-ol, benzyl cyanide, 2-methylisoborneol, geosmin (1-10-dimethyl-*trans*-9-decalol), and/or terpenes also being present. The fact that 1-butanol was detected at low levels without the other above mentioned MVOCs would indicate that their presence on the 2nd Floor was most likely not fungal growth related and attributable to personal products such as perfumes and other personal cosmetic products. All such data are well below the applicable Cal-OSHA 8-hour TWA PELs as defined in T8, CCR § 5155.

4.7 Airborne Volatile Organic Compounds

With the use of a direct-reading photoionization detector, VOCs were detected at a peak level of 0.3 ppm. Because these data were recorded at stationary locations at approximate breathing zone height, the results are expected to represent building occupant *exposure* potentials for those persons occupying or passing through the areas monitored. These data were well below the surrogate Cal-OSHA PELs that are often used for comparative purposes regarding VOC exposures, such as those for gasoline, hexane, and varnish makers and painters (VM&P) naphtha.



4.0 DISCUSSION (CONTINUED)

4.8 Airborne Ozone

O₃ was not detected at or above the Dräger instrument detection limits of 0.05 ppm.

4.9 Airborne Carbon Dioxide

On April 15, 2008, the direct-reading results indicated that CO₂ was detected at levels ranging from 454 and 542 ppm on the 2nd Floor. While these data were somewhat higher than the expected outdoor CO₂ levels, which generally range between 320 and 350 ppm, they are considered normal for occupied indoor environments and they are all well below the Cal-OSHA 8-hour TWA PEL for CO₂ of 5000 ppm (T8, CCR, § 5155). They are also below the level of 1000 ppm, which is essentially equivalent to the recommended upper limit for building occupant comfort and odor control established by ASHRAE (not greater than 700 ppm above the outdoor CO₂ value) as stated in ASHRAE 62-2001.

Based on historic studies performed by HygieneTech, building occupant complaints of "stuffy" air often begin when CO₂ levels exceed 800 ppm. HygieneTech has also found that some sensitive persons may experience discomfort, including eye irritation and headache, when CO₂ levels reach 1,000 ppm. Such symptoms are not believed to be the result of an unhealthful exposure to CO₂; rather, they are thought to be the result of exposure to other common indoor air pollutants which, if not exhausted and/or diluted, can accumulate over time.

4.10 Air Temperature and Relative Humidity

On April 15, 2008, air temperatures indoors ranged between 72.5 and 74.5 degrees Fahrenheit (°F). Based on the experience of HygieneTech, the air temperatures perceived as comfortable by most persons in office environments, and recommended by ASHRAE for occupant comfort, range between 68.0 and 74.5°F (winter) and 73.0 and 79.0°F (summer). The air temperatures recorded in the surveyed areas were within the comfort range recommended for the winter months.

Relative humidity data were recorded indoors at levels ranging from 23.4 to 24.0 percent. Such levels were well within the 20 to 60 percent relative humidity level range recommended by ASHRAE for occupant comfort. Note that HygieneTech recommends that the relative humidity in buildings not exceed 50 percent in order to limit the potential for fungal growth.

5.0 CONCLUSIONS

5.1 The airborne total and viable fungi data recorded in the surveyed areas showed airborne fungi levels that were generally below those recorded outdoors and therefore considered unremarkable. These data are not believed to pose a health risk beyond that posed by the outdoor environment where exposures to airborne fungi are expected. However, note that the airborne fungi results recorded within the ceiling plenum indicated somewhat higher levels of *Cladosporium* at two locations, the airborne data recorded within the occupied space would suggest that the conditions in the ceiling plenums did not appear to have adversely affected the indoor air quality on the 2ND Floor.



5.0 CONCLUSIONS (CONTINUED)

- 5.2 The surface assessment data involving the samples collected from various cubicle partitions and other surfaces throughout the 2ND Floor indicated no evidence of fungal growth or above-background levels of loose fungal spores on those surfaces. Additionally, the surface assessment data from the samples collected from the HVAC supply air registers throughout the floor indicated loose fungal spores involving *Alternaria*, *Cladosporium*, *Penicillium/Aspergillus* types, and/or *Ulocladium* on six of the eight registers sampled. Be advised that visible accumulation of debris, dust, and other particulates was observed on the reverse side of all sampled HVAC supply air registers, and that such conditions are indicative of an environment that may promote fungal growth. However, note that the airborne fungi results discussed above would suggest that such conditions did not appear to have adversely affected the indoor air quality on the 2ND Floor.
- 5.3 The airborne total and fibrous dust, VOC, and O₃, and CO₂ levels recorded during the survey were unremarkable. Collectively, the data were well below applicable Cal-OSHA 8-hour TWA PELs and/or other occupational, non-occupational, ASHRAE, or foreign guidelines. The data are not expected to represent conditions that pose a measurable health risk to the building occupants.
- The airborne MVOC data indicated the presence of 1-butanol at levels ranging from 525 ng/m³ to 548 ng/m³. Microbial growth related VOCs would not be expected to be present indoors without additional MVOCs such as ethanol, 1-octen-3-ol, 2-octen-1-ol, benzyl cyanide, 2-methylisoborneol, geosmin (1-10-dimethyl-*trans*-9-decalol), and/or terpenes also being present. The fact that 1-butanol was detected at low levels without the other above mentioned MVOCs would indicate that their presence on the 2nd Floor was most likely not fungal growth related and attributable to personal products such as perfumes and other personal cosmetic products. All such data are well below the applicable Cal-OSHA 8-hour TWA PELs as defined in T8, CCR § 5155.
- Indoor air temperatures ranged between 72.5 and 74.5 degrees Fahrenheit (°F) on the survey date. Based on the experience of HygieneTech, the air temperatures perceived as comfortable by most persons in office environments, and recommended by ASHRAE for occupant comfort, range between 68.0 and 74.5°F (winter) and 73.0 and 79.0°F (summer). The air temperatures recorded in the surveyed areas were within the comfort range recommended for the winter months. Relative humidity data were recorded indoors at levels ranging from 23.4 to 24.0 percent, levels that were well within the 20 to 60 percent relative humidity level range recommended by ASHRAE for occupant comfort. Note that HygieneTech recommends that the humidity in buildings not exceed 50 percent in order to limit the potential fungal growth.
- 5.6 Be advised that the data provided in this report only represent fungal growth and exposure potentials that existed at the time the survey was performed and at the precise sample locations only, the latter of which were selected based on the available background information provided. Note that fungal growth and exposure potentials may change due to changes in environmental conditions (such as those caused by water intrusion), use of mechanical systems, or other factors. Also be advised that additional fungal growth may exist at one or more locations in the structure that were not specifically assessed during the survey.



6.0 RECOMMENDATIONS

All such recommendations are based strictly on the assessment information and analytical data that were available to HygieneTech at the time this report was prepared. Be advised that, in order to establish data that accurately reflects all the fungal growth sites on the 2ND Floor, additional assessment evaluations may be required as more information is known regarding the history of water intrusion episodes in discrete building areas.

- 6.1 If not yet established, an accurate record of all air monitoring results should be maintained in accordance with Cal-OSHA regulation found in T8, CCR § 3204. All affected employees should be informed that the *exposure potential* data in this report exist and that those persons, or their representatives, have a right to access relevant exposure data and medical records.
- 6.2 Routine cleaning of the HVAC supply air registers on the 2ND Floor should be performed to preclude the build-up of dust and debris, which may potentially contribute to fungal growth on those surfaces.
- 6.3 A detail inspection of the ceiling plenum should be conducted to determine possible sources of water intrusion and/or fungal growth reservoirs.
- Also be advised that the exposure data recorded during the survey may not be sufficiently broad to adequately assess the suitability of the indoor air quality for all individuals, particularly those who are extremely sensitive to certain chemical and/or biological substances or for those individuals with immune system deficiencies. Although not expected, if persons occupying or passing through the 2nd Floor do experience non-specific ill effects of unknown etiology, then those affected should be referred to a medical professional in order to determine or specify the possible cause(s) of such reactions. If more information becomes available, further investigation and air monitoring may be warranted.

HYGIENE TECHNOLOGIES INTERNATIONAL, INC.

Brian P. Daly, CIH, PE

President

lawy >	Date:	August 8, 2008	
Kenny K. Hsi, CIH Technical Director			
Sien P. alely	Date:	August 8, 2008	
10000			

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 1

Results reported in spores per cubic meter of air (spores/M³)

Resu	ılts reported in spores			
SAMPLE NUMBER	20804001-TM01OUTME	20804001-TM02ME	20804001-TM03ME	20804001-TM04ME
SAMPLING LOCATION/ACTIVITIES	Outdoor; about 25 feet east of building; approximately five feet above ground/Normal outdoor activities	About six feet east of Column J23 area; approximately five feet above floor/Normal office activities	Column K23 area; Cubicle 043; about center; approximately five feet above floor/Normal office activities	Column K22 area; Cubicle 30; about center; approximately five feet above floor/Normal office activities
DATE	04-04-08	04-04-08	04-04-08	04-04-08
START/STOP	14:15:00/14:20:00	14:30:00/14:35:00	14:39:00/14:44:00	14:45:00/14:50:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria	Р			
Arthrinium	Р			
Ascospores	52		Р	
Aureobasidium				
Basidiospores	260	Р		100
Bipolaris/Drechslera group				
Botrytis				
Chaetomium	52			
Cladosporium	1,100	Р		Р
Curvularia				
Epicoccum				
Nigrospora				
Oidium	Р			
Penicillium/Aspergillus types	52	100	100	210
Pithomyces				
Rusts	260		Р	
Smuts (Periconia, Myxomycetes)	100			
Stachybotrys	100			
Stemphylium				
Torula	310		Р	
Ulocladium				
Unidentified mitosporic fungi	520		Р	100
Background particulates*	Light	Light	Light	Light
TOTAL**	2,800	100	100	410

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 2

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	Its reported in sport 20804001-TM05ME	20804001-TM06ME	20804001-TM07ME	20804001-TM08ME
SAMPLING LOCATION/ACTIVITIES	Column M22 area; Cubicle 58; about center; approximately five feet above floor/Normal office activities	Column N22 area; Cubicle 37; about center; approximately five feet above floor/Normal office activities	Area between Columns N21 and N22; Cubicle 103; about center; approximately five feet above floor/Normal office activities	About 15 feet north of Break Room 214; approximately five feet above floor/Normal office activities
DATE	04-04-08	04-04-08	04-04-08	04-04-08
START/STOP	14:54:00/14:59:00	15:00:00/15:05:00	15:09:00/15:14:00	15:15:00/15:20:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				
Arthrinium				
Ascospores				
Aureobasidium				
Basidiospores	52	100	160	
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium	52	52	Р	
Curvularia				
Epicoccum				
Myrothecium				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types	Р	Р		Р
Pithomyces				
Rusts				
Smuts (Periconia, Myxomycetes)		52		Р
Stachybotrys				
Torula				
Ulocladium				
Unidentified mitosporic fungi	Р	52	Р	Р
Background particulates*	Light	Light	Light	Light
TOTAL**	100	260	160	<52

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 3

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-TM09ME	20804001-TM10ME	20804001-TM11ME	20804001-TM12ME
SAMPLING LOCATION/ACTIVITIES	Area between Column N19 and N20 area; Cubicle 134; about center; approximately five feet above floor/Normal office activities	Column N19 area; Cubicle 119; about center; approximately five feet above floor/Normal office activities	Column N18 area; Cubicle 127; about center; approximately five feet above floor/Normal office activities	Column L18 area; Cubicle 007; about center; approximately five feet above floor/Normal office activities
DATE	04-04-08	04-04-08	04-04-08	04-04-08
START/STOP	15:24:00/15:29:00	15:30:00/15:35:00	15:36:00/15:41:00	15:45:00/15:50:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				
Arthrinium				
Ascospores		Р	52	
Aureobasidium				
Basidiospores		52	Р	Р
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium	Р	52	52	52
Curvularia				
Epicoccum				
Myrothecium				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types	52		160	52
Pithomyces				
Rusts				
Smuts (Periconia, Myxomycetes)			Р	
Stachybotrys				
Torula			Р	
Ulocladium				
Unidentified mitosporic fungi			Р	52
Background particulates*	Very light	Moderate	Moderate	Moderate
TOTAL**	52	100	260	160

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 4

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-TM13ME	20804001-TM14ME	20804001-TM15ME	20804001-TM16ME
SAMPLING LOCATION/ACTIVITIES	Column K18 area; Cubicle 11; about center; approximately five feet above floor/Normal office activities	Column K19 area; Cubicle 15; about center; approximately five feet above floor/Normal office activities	About two feet east of Column K19; approximately five feet above floor/Normal office activities	Column J19 area; Cubicle 23; about center; approximately five feet above floor/Normal office activities
DATE	04-04-08	04-04-08	04-04-08	04-04-08
START/STOP	15:54:00/15:59	16:00:00/16:05:00	16:06:00/16:11:00	16:15:00/16:20:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				
Arthrinium				
Ascospores			52	
Aureobasidium				
Basidiospores		52	Р	Р
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium	52		100	
Curvularia				
Epicoccum				
Myrothecium				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types	52	100	210	
Pithomyces				
Rusts				
Smuts (Periconia, Myxomycetes)	52			Р
Stachybotrys				
Torula				
Ulocladium				
Unidentified mitosporic fungi		100	Р	52
Background particulates*	Moderate	Moderate	Moderate	Very light
TOTAL**	160	250	360	52

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 5

Results reported in spores per cubic meter of air (spores/M³)

Resu	ults reported in spo	res per cubic meter of	f air (spores/M³)	
SAMPLE NUMBER	20804001-TM17ME	20804001-TM18OUTME	20804001-TM01CCJL	20804001-TM02CCJL
SAMPLING LOCATION/ACTIVITIES	Room 208; about center; approximately five feet above floor/Normal office activities	Outdoors; about 30 feet east of building; approximately five feet above ground/Normal outdoor activities	Area between Column K23 and J23 area; approximately five feet east of Cubicle 81; within ceiling plenum/Sampling activities only	Column L23 area; approximately three feet south of Cubicle 55; within ceiling plenum/Sampling activities only
DATE	04-04-08	04-04-08	04-04-08	04-04-08
START/STOP	16:21:00/16:26:00	16:35:00/16:40:00	13:36:00/13:41:00	13:48:00/ 13:53:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				100
Arthrinium				
Ascospores		52	Р	160
Aureobasidium				
Basidiospores		990	52	680
Bipolaris/Drechslera group		52		
Botrytis				Р
Chaetomium		52		
Cladosporium		1,600	210	3,000
Curvularia				
Epicoccum		Р		
Fusarium			Р	
Nigrospora				
Oidium		52	52	
Other brown				260
Penicillium/Aspergillus types		Р	160	
Pithomyces				
Rusts		Р		Р
Smuts (Periconia, Myxomycetes)	Р	310	52	260
Stachybotrys		Р	Р	
Torula				52
Ulocladium				Р
Unidentified mitosporic fungi		310	Р	Р
Background particulates*	Very light	Light	Moderate	Heavy
TOTAL**	<52	3,400	530	4,500

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 6

Results reported in spores per cubic meter of air (spores/M³)

SAMPLING LOCATION/ACTIVITIES Column N22 area; Cubicle 37; about center; within ceiling plenum/Sampling activities only DATE Column N20 area; about two feet west of Cubicle 110; within ceiling plenum/Sampling activities only Od-04-08 Column N20 area; about two feet west of Cubicle 120; within ceiling plenum/Sampling activities only Od-04-08 Column N19 area; approximately two feet east of Cubicle 120; within ceiling plenum/Sampling activities only Od-04-08 Od-04-08 Od-04-08 Od-04-08 Od-04-08		Results reported in spores per cubic meter of air (spores/M³)				
LOCATION/ACTIVITIES Cubicle 37; about center: within celling plenum/Sampling activities only lenum/Sampling activities only about two feet west of Cubicle 110; within celling plenum/Sampling activities only Cubicle 110; within celling plenum/Sampling plenum/Sampling activities only Subject of Cubicle 120; within celling plenum/Sampling activities only Cubicle 110; within celling plenum/Sampling activities only Activation of Cubicle 120; within celling plenum/Sampling activities only Activation of Cubicle 120; within celling plenum/Sampling activities only Activation of Cubicle 120; within celling plenum/Sampling activities only Activation of Cubicle 120; within celling plenum/Sampling activities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling plenum/Sampling pactivities only Activation of Cubicle 120; within celling pactivities only Activation	SAMPLE NUMBER	20804001-TM03CCJL	20804001-TM04CCJL	20804001-TM05CCJL	20804001-TM06CCJL	
START/STOP 13:58:00/14:03:00 14:08:00/14:13:00 14:18:00/14:23:00 14:25:00/14:30:00 SAMPLE TIME 5 minutes 5 minutes 5 minutes 5 minutes Alternaria 100 P 4 4 Arthrinium		Cubicle 37; about center; within ceiling plenum/Sampling	about two feet west of Cubicle 110; within ceiling plenum/Sampling activities only	approximately two feet east of Cubicle 120; within ceiling plenum/Sampling	Column L18 area; Cubicle 007; about center; within ceiling plenum/Sampling activities only	
SAMPLE TIME 5 minutes 5 minutes 5 minutes Alternaria 100 P ————————————————————————————————————	DATE	04-04-08	04-04-08	04-04-08	04-04-08	
Alternaria 100 P	START/STOP	13:58:00/14:03:00	14:08:00/14:13:00	14:18:00/14:23:00	14:25:00/14:30:00	
Arthrinium S2 52 52 52 52 Aureobasidium S2 52 52 52 S2 S2 </td <td>SAMPLE TIME</td> <td>5 minutes</td> <td>5 minutes</td> <td>5 minutes</td> <td>5 minutes</td>	SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes	
Ascospores 52 52 52 Aureobasidium 100 260 52 Basidiospores 470 100 260 52 Bipolaris/Drechslera group 100 100 100 100 Botrytis P 100 160 160 52 100 160 160 52 100	Alternaria	100	Р			
Aureobasidium 470 100 260 52 Basidiospores 470 100 260 52 Bipolaris/Drechslera group	Arthrinium					
Basidiospores 470 100 260 52 Bipolaris/Drechslera group Bipolaris/Drechslera group Bipolaris/Drechslera group Bipolaris/Drechslera group Botrytis P Standard Green Gre	Ascospores	52	52	52		
Bipolaris/Drechslera group Botrytis P	Aureobasidium					
P	Basidiospores	470	100	260	52	
Chaetomium 2,000 160 160 52 Curvularia	Bipolaris/Drechslera group					
Cladosporium 2,000 160 160 52 Curvularia	Botrytis	Р				
Curvularia Fusarium Epicoccum P Nigrospora P Oidium P Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) 210 P Stachybotrys Stemphylium P Torula Ulocladium P Unidentified mitosporic fungi P Unidentified zygomycetes Background particulates* Moderate Moderate	Chaetomium					
Fusarium Epicoccum P Nigrospora P Oidium P Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) Stemphylium P Torula Ulocladium P Unidentified mitosporic fungi P Moderate Moderate Moderate	Cladosporium	2,000	160	160	52	
Epicoccum P Nigrospora P Oidium P Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) 210 P 100 P Stachybotrys Stemphylium P Torula P Ulocladium P Unidentified mitosporic fungi 210 160 P Sackground particulates* Moderate Moderate Moderate	Curvularia					
Nigrospora P Oidium P Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) 210 P 100 P Stachybotrys Stemphylium P Torula P Unidentified mitosporic fungi 210 160 P Unidentified zygomycetes P Background particulates* Moderate Moderate Moderate	Fusarium					
Oidium P Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) 210 P 100 P Stachybotrys Stemphylium P Torula P Unidentified mitosporic fungi 210 160 P Unidentified zygomycetes P Background particulates* Moderate Moderate Moderate	Epicoccum			Р		
Penicillium/Aspergillus types 160 260 100 100 Pithomyces Rusts P Smuts (Periconia, Myxomycetes) 210 P 100 P Stachybotrys Stemphylium P Torula 52 Ulocladium P Unidentified mitosporic fungi 210 160 P Background particulates* Moderate Moderate Moderate	Nigrospora	Р				
Pithomyces Rusts P Smuts (Periconia, Myxomycetes) Stachybotrys Stemphylium P Torula Ulocladium P Unidentified mitosporic fungi Background particulates* Moderate P Moderate Moderate P Moderate Moderate	Oidium	Р				
Rusts P 100 P 100 P Stachybotrys P 52 Ulocladium P 52 Unidentified zygomycetes P 52 Background particulates* Moderate Moderate Moderate	Penicillium/Aspergillus types	160	260	100	100	
Smuts (Periconia, Myxomycetes) Stachybotrys Stemphylium P Torula Ulocladium P Unidentified mitosporic fungi Dackground particulates* Moderate P 100 P 100 P 100 P 100 P 100 P 52 P 52 Moderate Moderate Moderate Moderate	Pithomyces					
Stachybotrys P Stemphylium P Torula 52 Ulocladium P Unidentified mitosporic fungi 210 160 P Unidentified zygomycetes P 52 Background particulates* Moderate Moderate Moderate	Rusts	Р				
Stemphylium P Torula 52 Ulocladium P Unidentified mitosporic fungi 210 Unidentified zygomycetes P Background particulates* Moderate Moderate Moderate	Smuts (Periconia, Myxomycetes)	210	Р	100	Р	
Torula 52 Ulocladium P Unidentified mitosporic fungi 210 160 P Unidentified zygomycetes P 52 Background particulates* Moderate Moderate Moderate Moderate	Stachybotrys					
Ulocladium P	Stemphylium	Р				
Unidentified mitosporic fungi 210 160 P Unidentified zygomycetes P 52 Background particulates* Moderate Moderate Moderate	Torula			52		
Unidentified zygomycetes P 52 Background particulates* Moderate Moderate Moderate Moderate	Ulocladium	Р				
Background particulates* Moderate Moderate Moderate Moderate	Unidentified mitosporic fungi	210	160		Р	
•	Unidentified zygomycetes		Р	52		
TOTAL** 3,200 730 780 200	Background particulates*	Moderate	Moderate	Moderate	Moderate	
	TOTAL**	3,200	730	780	200	

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138 **AIRBORNE TOTAL FUNGI RESULTS** 2ND FLOOR **SACRAMENTO, CALIFORNIA APRIL 4, 14, AND 15, 2008**

Page 7

Results reported in spores per cubic meter of air (spores/M3)

SAMPLE NUMBER	20804001-TM07CCJL	20804001-TM08CCJL	20804001-TM17OUTJL	20804001-TM18JL
SAMPLING LOCATION/ACTIVITIES	Column K19 area; approximately three feet east of Cubicle 20; within ceiling plenum/Sampling activities only	Room 208; about center; within ceiling plenum/Sampling activities only	Outdoor; about 20 feet east of building; approximately five feet above ground/Normal outdoor activities	Room 208; about center; approximately five feet above floor/Sampling activities only
DATE	04-04-08	04-04-08	04-14-08	04-14-08
START/STOP	14:32:00/14:37:00	14:41:00/14:46:00	13:18:00/13:23:00	13:40:00/13:45:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria	Р		27	
Arthrinium				
Ascospores	Р		53	
Aureobasidium				
Basidiospores	260	100	320	
Bipolaris/Drechslera group				
Botrytis		Р		
Chaetomium				
Cladosporium	100	100	480	
Curvularia				
Fusarium				
Epicoccum				
Nigrospora				
Oidium	Р		13	53
Penicillium/Aspergillus types	360		427	
Pithomyces				
Rusts	Р		13	
Smuts (Periconia, Myxomycetes)	100		120	
Stachybotrys				
Stemphylium				
Torula				
Ulocladium				
Unidentified mitosporic fungi	Р	Р		
Hyphal fragments			360	<13
Background particulates*	Moderate	Light	3+	1+
TOTAL**	820	200	1,453	53

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138 **AIRBORNE TOTAL FUNGI RESULTS** 2ND FLOOR **SACRAMENTO, CALIFORNIA APRIL 4, 14, AND 15, 2008**

Results reported in spores per cubic meter of air (spores/M³)					
SAMPLE NUMBER	20804001-TM19JL	20804001-TM20JL	20804001-TM21JL	20804001-TM22JL	
SAMPLING LOCATION/ACTIVITIES	Column K22 area; Cubicle 31; about center; approximately five feet above floor/Normal office activities	Area between Columns K23 and J23; Cubicle 81; about center; approximately five feet above floor/Normal office activities	Room 207; about center; approximately five feet above floor/Normal office activities	Column L23 area; Cubicle 55; about center; approximately five feet above floor/Normal office activities	
DATE	04-14-08	04-14-08	04-14-08	04-14-08	
START/STOP	13:45:00/13:50:00	13:51:00/13:56:00	13:57:00/14:02:00	14:04:00/14:09:00	
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes	
Alternaria					
Arthrinium					
Ascospores					
Aureobasidium					
Basidiospores					
Bipolaris/Drechslera group					
Botrytis					
Chaetomium					
Cladosporium				53	
Curvularia					
Fusarium					
Epicoccum					
Nigrospora					
Oidium					
Penicillium/Aspergillus types	53	53			
Pithomyces					
Rusts					
Smuts (Periconia, Myxomycetes)			27		
Stachybotrys					
Stemphylium					
Torula					
Ulocladium					
Unidentified mitosporic fungi					
Hyphal fragments	<13	<13	<13	<13	
Background particulates*	1+	1+	1+	1+	
TOTAL**	53	53	27	53	

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 9

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	Its reported in sport	20804001-TM24JL	20804001-TM25JL	20804001-TM26JL
SAMPLING LOCATION/ACTIVITIES	Column M23 area; Cubicle 51; about center; approximately five feet above floor/Normal office activities	Column N22 area; Cubicle 37; about center; approximately five feet above floor/Normal office activities	Area between Columns N21 and N22; Cubicle 103; about center; approximately five feet above floor/Normal office activities	Column N21 area; Cubicle 106; about center; approximately five feet above floor/Normal office activities
DATE	04-14-08	04-14-08	04-14-08	04-14-08
START/STOP	14:10:00/14:15:00	14:16:00/14:21:00	14:22:00/14:27:00	14:28:00/14:33:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				
Arthrinium				
Ascospores				
Aureobasidium				
Basidiospores		53		53
Bipolaris/Drechslera group				
Botrytis				
Chaetomium	13			
Cladosporium		53	53	
Curvularia				
Epicoccum				
Myrothecium				
Nigrospora				
Oidium		13		
Other brown				
Penicillium/Aspergillus types	53		107	53
Pithomyces				
Rusts				
Smuts (Periconia, Myxomycetes)				
Stachybotrys				
Torula				
Ulocladium				
Hyphal fragments	<13	<13	<13	<13
Background particulates*	1+	2+	2+	2+
TOTAL**	66	119	160	106

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 10

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-TM27JL	20804001-TM28JL	20804001-TM29JL	20804001-TM30JL
SAMPLING LOCATION/ACTIVITIES	Area between Columns N20 and N21; Cubicle 138; about center; approximately five feet above floor/Normal office activities	Column N20 area; Cubicle 110; about center; approximately five feet above floor/Normal office activities	Column M18 area; Cubicle 124; about center; approximately five feet above floor/Normal office activities	Column M18 area; Cubicle 005; about center; approximately five feet above floor/Normal office activities
DATE	04-14-08	04-14-08	04-14-08	04-14-08
START/STOP	14:33:00/14:38:00	14:39:00/14:44:00	14:45:00/14:50:00	15:11:00/15:16:00
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes
Alternaria				
Arthrinium				
Ascospores				
Aureobasidium				
Basidiospores	53			
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium	53	53	107	160
Curvularia				
Epicoccum				
Myrothecium				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types	880	293	53	
Pithomyces				
Rusts	13			
Smuts (Periconia, Myxomycetes)	67			27
Stachybotrys				
Torula				
Ulocladium				
Hyphal fragments	<13	<13	<13	<13
Background particulates*	2+	2+	1+	1+
TOTAL**	1,066	346	160	187

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 11

Results reported in spores per cubic meter of air (spores/M³)

Resu	Results reported in spores per cubic meter of air (spores/M³)				
SAMPLE NUMBER	20804001-TM31JL	20804001-TM32JL	20804001-TM33JL	20804001-TM34OUTJL	
SAMPLING LOCATION/ACTIVITIES	Column K18 area; Cubicle 011; about center; approximately five feet above floor/Normal office activities	Area between Columns K19 and K20; Cubicle 020; about center; approximately five feet above floor/Normal office activities	Column J19 area; approximately five feet east of Cubicle 024; approximately five feet above floor/Normal office activities	Outdoors; approximately 20 feet east of building; approximately five feet above ground/Normal office activities	
DATE	04-14-08	04-14-08	04-14-08	04-14-08	
START/STOP	15:17:00/15:22:00	15:24:00/15:29:00	14:30:00/14:35:00	15:44:00/15:49:00	
SAMPLE TIME	5 minutes	5 minutes	5 minutes	5 minutes	
Alternaria				13	
Arthrinium					
Ascospores					
Aureobasidium					
Basidiospores		53	160	267	
Bipolaris/Drechslera group					
Botrytis					
Chaetomium					
Cladosporium			53	2,130	
Curvularia				13	
Epicoccum					
Myrothecium					
Nigrospora					
Oidium				13	
Other brown					
Penicillium/Aspergillus types		53	53	747	
Pithomyces					
Rusts				13	
Smuts (Periconia, Myxomycetes)		13		200	
Stachybotrys				27	
Torula				40	
Ulocladium					
Hyphal fragments	<13	13	<13	120	
Background particulates*	1+	2+	1+	3+	
TOTAL**	<13	119	266	3,463	

P = Spores present

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-138
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 14, AND 15, 2008

Page 12

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-TM35JL	es per cubic meter of 20804001-TM36JL	20804001-TM37JL	
SAMPLING LOCATION/ACTIVITIES	Column K22 area; Cubicle 036; southeastern corner; approximately five feet above floor/Normal office activities	Column M23 area; Cubicle 056; northwest corner; approximately five feet above floor/Normal office activities	Column M22 area; Cubicle 100; about center; approximately five feet above floor/Normal office activities	This Column intentionally left blank
DATE	04-15-08	04-15-08	04-15-08	
START/STOP	11:17:00/11:22:00	11:25:00/11:30:00	11:31:00/11:36:00	
SAMPLE TIME	5 minutes	5 minutes	5 minutes	
Alternaria				
Arthrinium				
Ascospores				
Aureobasidium				
Basidiospores	160		Р	
Bipolaris/Drechslera group				
Botrytis				
Chaetomium				
Cladosporium		210	100	
Curvularia				
Epicoccum				
Nigrospora				
Oidium				
Other brown				
Penicillium/Aspergillus types	Р			
Pithomyces				
Rusts				
Smuts (Periconia, Myxomycetes)	Р	Р	Р	
Stachybotrys				
Torula				
Ulocladium				
Unidentified mitosporic fungi	Р	52	Р	
Hyphal fragments				
Background particulates*	Moderate	Moderate	Light	
TOTAL**	160	260	100	

^{*}Background particulates is an indication of the amount of non-biological particulate matter present on the media and is graded (from least to greatest) as very light, light, moderate, heavy and very heavy or as 1+ to 4+.

^{**} Note that all reported counts have been rounded to no more than two significant figures based on the sampling and analytical methods used, and therefore the total count may not equal the sum of the individual counts in a column.

CLIENT: California State Board of Equalization

450 N Street Sacramento, California 94279 **APPENDIX A**

TABLE 20803001-139
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 14, 2008

Page 1

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-VM01OUTJL	20804001-VM02JL	20804001-VM03JL	20804001-VM04JL
SAMPLING LOCATION/ACTIVITIES	Outdoor; about 20 feet east of building; approximately five feet above ground/Normal outdoor activities	Room 208; about center; approximately five feet above floor/Sampling activities only	Column J23 area; Cubicle 81; about center; approximately five feet above floor/Normal office activities	Column L23 area; Cubicle 55; about center; approximately five feet above floor/Normal office activities
START/STOP	13:20:00/13:22:00	13:42:00/13:44:00	13:53:00/13:55:00	14:07:00/14:09:00
SAMPLE TIME	2 minutes	2 minutes	2 minutes	2 minutes
Acremonium				
Alternaria	18			
Aphanocladium				
Arthrospore-former		18		
Aspergillus fumigatus				
Aspergillus nidulans				
Aspergillus niger				
Aspergillus versicolor				
Aureobasidium				
Botrytis				
Chaetomium	18			
Cladosporium	1,150		18	18
Curvularia				
Epicoccum				
Nigrospora				
Memnoniella				
Myrothecium				
Non-sporulating fungi	35			
Others				
Paecilomyces				
Penicillium				
Stachybotrys				
Torula herbarum				
Trichoderma				
Ulocladium				
Yeasts	18			
TOTAL	1,239	18	18	18

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-139
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 14, 2008

Page 2

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-VM05JL	20804001-VM06JL	20804001-VM07JL	20804001-VM08JL
SAMPLING LOCATION/ACTIVITIES	Column N22 area; Cubicle 037; about center; approximately five feet above floor/Normal office activities	Column N21 area; Cubicle 106; about center; approximately five feet above floor/Normal office activities	Column N20 area; Cubicle 110; about center; approximately five feet above floor/Normal office activities	Column M18 area; Cubicle 005; about center; approximately five feet above floor/Normal office activities
START/STOP	14:18:00/14:20:00	14:30:00/14:32:00	14:41:00/14:43:00	15:13:00/15:15:00
SAMPLE TIME	2 minutes	2 minutes	2 minutes	2 minutes
Acremonium				
Alternaria			18	
Aphanocladium				
Arthrospore former				
Aspergillus fumigatus				
Aspergillus nidulans				
Aspergillus niger				
Aspergillus versicolor				
Aureobasidium				
Botrytis				
Chaetomium				
Cladosporium			35	18
Curvularia				
Epicoccum				
Nigrospora				
Memnoniella				
Myrothecium				
Non-sporulating fungi				
Others				
Paecilomyces				
Penicillium	18	35		
Stachybotrys				
Torula herbarum				
Trichoderma				
Ulocladium				
Yeasts		18		
TOTAL	18	53	53	18

CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

APPENDIX A

TABLE 20803001-139
AIRBORNE TOTAL FUNGI RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 14, 2008

Page 3

Results reported in spores per cubic meter of air (spores/M³)

SAMPLE NUMBER	20804001-VM09JL	20804001-VM10OUTJL	,	
SAMPLING LOCATION/ACTIVITIES	Area between Columns K19 and K20; Cubicle 20; about center; approximately five feet above floor/Normal office activities	Outdoors; about 20 feet east of building; approximately five feet above ground/Normal outdoor activities	This column intentionally left blank	This column intentionally left blank
START/STOP	15:25:00/15:27:00	15:47:00/15:49:00		
SAMPLE TIME	2 minutes	2 minutes		
Acremonium				
Alternaria		18		
Aphanocladium		18		
Arthrospore former				
Aspergillus fumigatus	18			
Aspergillus nidulans		18		
Aspergillus niger		18		
Aspergillus versicolor				
Aureobasidium		35		
Botrytis				
Chaetomium				
Cladosporium	18	1,380		
Curvularia				
Epicoccum				
Nigrospora				
Memnoniella				
Myrothecium				
Non-sporulating fungi	18			
Others				
Paecilomyces				
Penicillium	18	53		
Stachybotrys				
Torula herbarum				
Trichoderma				
Ulocladium				
Yeasts	18	53		
TOTAL	90	1,593		



CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

TABLE 20803001-140
SURFACE FUNGAL GROWTH POTENTIALS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

						rayeı
SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- TL01JL	Column K22 area; Cubicle 30; eastern cubicle partition; about center; from top horizontal surface	Light dander Very light fibers Very light particulates	Trace	None	Trace Epicoccum ¹	Background
20804001- TL02JL	Column K22 area; Cubicle 36; western cubicle partition; about center; from top horizontal surface	Moderate dander Moderate fibers Moderate particulates Very light insect parts	Trace	None	Trace Cladosporium Trace Curvularia Trace Epicoccum Trace Oidium ¹	Background
20804001- TL03JL	Area between Column K23 and J23; Cubicle 81; northern cubicle partition; about center; from top horizontal surface	Light dander Light particulates Very light fibers	Trace	None	None	Background
20804001- TL04JL	Room 207; eastern cubicle partition cabinetry; about center; from top horizontal surface	Light dander Light fibers Very light particulates	Trace	None	None	Background
20804001- TL05JL	Column M22 area; Cubicle 57; southern cubicle partition; about center; from top horizontal surface	Light dander Light particulates Very light fibers	Trace	None	Trace Cladosporium	Background

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.



CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

TABLE 20803001-140
SURFACE FUNGAL GROWTH POTENTIALS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

						ı aye z
SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- TL06JL	Column N22 area; Cubicle 042; western cubicle partition; about center; from top horizontal surface	Light dander Very light fibers Very light particulates	Trace	None	None	Background
20804001- TL07JL	Column N22 area; Cubicle 143; southern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	None	None	None	Background
20804001- TL08JL	Area between Columns N21 and N22; Cubicle 103; western cubicle partition; about center; from top horizontal surface	Light dander Very light fibers Very light particulates	Trace	None	None	Background
20804001- TL09JL	Column N21 area; Cubicle 141; eastern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	Trace	None	Trace Penicillium/ Aspergillus types	Background
20804001- TL10JL	Column N21 area; Cubicle 139; eastern cubicle partition; about center; from top horizontal surface	Light dander Light fibers Light particulates	Trace	None	Trace Penicillium/ Aspergillus types	Background

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.



CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

SURFACE FUNGAL GROWTH POTENTIALS 2^{ND} FLOOR SACRAMENTO, CALIFORNIA

APRIL 4, 2008

TABLE 20803001-140

		1				
SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- TL11JL	Column N20 area; Cubicle 111; eastern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	Trace	None	Trace Penicillium/ Aspergillus types ¹	Background
20804001- TL12JL	Column N19 area; Cubicle 133; eastern cubicle partition; about center; from top horizontal surface	Moderate dander Light fibers Very light particulates	Trace	None	Trace Cladosporium Trace Penicillium/ Aspergillus types	Background
20804001- TL13JL	Area between Columns N18 and N19; Cubicle 122; eastern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	Trace	None	Trace Penicillium/ Aspergillus types ¹	Background
20804001- TL14JL	Column N18 area; Cubicle 001; southern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	None	None	None	Background
20804001- TL15JL	Area between Columns L18 and M18; Cubicle 006; northern cubicle partition; about center; from top horizontal surface of plastic	Very light dander Very light fibers Very light particulates	Trace	None	Trace Cladosporium ¹	Background

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.



CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

TABLE 20803001-140
SURFACE FUNGAL GROWTH POTENTIALS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

					1	ı aye 4
SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- TL16JL	Column K18 area; Cubicle 011; southern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	Trace	None	None	Background
20804001- TL17JL	Column K18 area; Cubicle 016; northern cubicle partition; about center; from top horizontal surface	Very light dander Very light fibers Very light particulates	Trace	None	Trace Cladosporium ¹	Background
20804001- TL18JL	Area between Columns K19 and K20; Cubicle 20; southern cubicle partition; about center; from top horizontal surface	Light dander Light particulates Very light fibers	Trace	None	Trace Penicillium/ Aspergillus types	Background
20804001- TL19JL	Area between Columns K19 and K20; Cubicle 26; southern cubicle partition; about center; from top horizontal surface	Light dander Light particulates Light fibers	Trace	None	Trace Cladosporium Trace Penicillium/ Aspergillus types Trace unidentified mitosporic fungi	Background

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.



CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

TABLE 20803001-140
SURFACE FUNGAL GROWTH POTENTIALS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- TL20JL	Room 208; Computer 17; monitor; about center; from top horizontal surface	Moderate dander Moderate fibers Light particulates	Trace	None	Trace Bipolaris/ Drechslera group ¹ Trace Cladosporium Trace Penicillium/ Aspergillus types Trace Torula ¹ Trace Ulocladium ¹	Background
20804001- S01JL	Column K23 area; about five feet east of Cubicle 81; ceiling; from reverse side of HVAC supply air register	Moderate	Few	None	None	Background
20804001- S02JL	Column L23 area; approximately three feet south of Cubicle 55; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many Ulocladium spores detected	Possible settling from unknown sources
20804001- S03JL	Column N22 area; Cubicle 37; about center; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many Ulocladium and Cladosporium spores detected	Possible settling from unknown sources
20804001- S04JL	Column N20 area; about two feet west of Cubicle 110; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many Ulocladium and colorless spores typical of Penicillium/ Aspergillus detected	Possible settling from unknown sources

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.



CLIENT: California State Board of Equalization 450 N Street

Sacramento, California 94279

TABLE 20803001-140
SURFACE FUNGAL GROWTH POTENTIALS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

SAMPLE NUMBER	SAMPLING LOCATION	AMORPHOUS DEBRIS	MISCELLANEOUS FUNGI/POLLEN*	FUNGI SEEN WITH UNDERLYING MYCELIAL AND/OR SPORULATING STRUCTURES**	OTHER COMMENTS	GENERAL IMPRESSION
20804001- S05JL	Column N19 area; about two feet east of Cubicle 120; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many colorless spores typical of <i>Penicillium/</i> <i>Aspergillus</i> detected	Possible settling from unknown sources
20804001- S06JL	Column L18 area; Cubicle 007; about center; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many colorless spores typical of Penicillium/ Aspergillus detected	Possible settling from unknown sources
20804001- S07JL	Column K19 area; about three feet east of Cubicle 20; ceiling; from reverse side of HVAC supply air register	Moderate	Few	None	None	Background
20804001- S08JL	Room 208; about center; ceiling; from reverse side of HVAC supply air register	Heavy	Few	None	Many Alternaria spores detected	Possible settling from unknown sources

^{*}Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{**}Quantities of fungi are graded (from least to greatest) as <1+ to 4+ or as none, trace, few, numerous, and massive.

¹ Single spore observed.

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279 **APPENDIX A**



TABLE 20803001-141
AIRBORNE FIBERS RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
MARCH 28, 2008

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	START/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (f/cc)	PEL (f/cc)
Area Sample	Column K22 area; Cubicle 030; about center;	N/A	20803001-	9:02/	466	Fibers	0.008	0.1
	approximately six feet above floor/Normal office activities		F17JL	16:48	minutes			
Area Sample	Column K19 area; about two feet west of Cubicle 021;	N/A	20803001-	9:05/	472	Fibers	0.007	0.1
	approximately six feet above floor/Normal office activities		F18JL	16:57	minutes			
Area Sample	Room 208; about center; approximately six feet above	N/A	20803001-	9:08/	470	Fibers	0.012	0.1
	floor/Normal office activities		F19JL	16:58	minutes			
Area Sample	Column L18 area; about three feet west of Cubicle 007;	N/A	20803001-	9:09/	472	Fibers	0.011	0.1
	approximately six feet above floor/Normal office activities		F20JL	17:01	minutes			
Area Sample	Area between Columns N18 and N19; Cubicle 122; about	N/A	20803001-	9:11/	471	Fibers	0.008	0.1
	center; approximately six feet above floor/Normal office		F21JL	17:02	minutes			
	activities							
Area Sample	Area between Columns N20 and N21 area; Cubicle 109;	N/A	20803001-	9:14/	471	Fibers	0.007	0.1
	about center; approximately six feet above floor/Normal		F22JL	17:05	minutes			
	office activities							
Area Sample	Area between Columns M22 and M23 area; Cubicle 056;	N/A	20803001-	9:15/	471	Fibers	0.007	0.1
	about center; approximately five feet above floor/Normal		F23JL	17:06	minutes			
	office activities							
Blank	N/A	N/A	20803001-	N/A	N/A	Fibers	All data blank	N/A
			F24BLANKJL				corrected	

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

<: Less than

f/cc: Fibers per cubic centimeter of air

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

APPENDIX A



TABLE 20803001-142
AIRBORNE TOTAL DUST RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 4, 2008

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	STAR T/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (mg/M³)	PEL (mg/M³)
Area Sample	Column K23 area; about five feet south of Cubicle 043; approximately six feet above floor/Normal office activities	N/A	20804001- TD17JL	9:50/ 14:51	301 minutes	Total dust	<0.17	10
Area Sample	Column N22 area; about two feet north of Cubicle 037; approximately six feet above floor/Normal office activities	N/A	20804001- TD18JL	9:55/ 14:53	298 minutes	Total dust	<0.17	10
Area Sample	Area between Columns N20 and N21; about five feet south of Cubicle 138; approximately six feet above floor/Normal office activities	N/A	20804001- TD19JL	9:58/ 14:57	299 minutes	Total dust	<0.17	10
Area Sample	Area between Columns N18 and N19; about six feet south of Cubicle 130; ;approximately six feet above floor/Normal office activities	N/A	20804001- TD20JL	10:01/ 15:00	299 minutes	Total dust	<0.17	10
Area Sample	Column L18 area; about five feet west of Cubicle 007; approximately six feet above floor/Normal office activities	N/A	20804001- TD21JL	10:04/ 15:02	298 minutes	Total dust	<0.17	10
Area Sample	Area between Columns K19 and K20 area; about six feet east of Cubicle 026; approximately six feet above floor/Normal office activities	N/A	20804001- TD22JL	10:06/ 15:04	298 minutes	Total dust	<0.17	10
Area Sample	Room 208; about center; approximately six feet above floor/Normal office activities	N/A	20804001- TD23JL	10:09/ 15:06	297 minutes	Total dust	<0.17	10
Blank	N/A	N/A	20804001- TD24BLANKJL	N/A	N/A	Total dust	All data blank corrected	N/A

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

<: Less than

mg/M³: Milligrams per cubic meter

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

APPENDIX A

TABLE 20803001-143
MICROBIAL VOLATILE ORGANIC COMPOUNDS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 7, 2008

Page 1

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	START/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (mg/M³)	PEL (mg/M³)
Area Sample	Column K23 area; about three feet south of	N/A	20804001-	13:59/	95	3-Methylfuran	nd	N/A
	Cubicle 43; approximately six feet above		M01JL	15:34	minutes	2-Methyl-1-propanol	nd	N/A
	floor/Normal office activities					1-Butanol	548 x10 ⁻⁶	300
						3-Methyl-2-butanol	nd	N/A
						2-Pentanol	nd	N/A
						3-Methyl-2-butanol	nd	N/A
						Methyl disulfide	nd	N/A
						Ethyl isobutyrate	nd	N/A
						2-Hexanone	nd	410
						2-Heptanone	nd	468
						5-Methyl-3-heptanone	nd	N/A
						1-Octen-3-ol	nd	N/A
						3-Octanone	nd	N/A
						3-Octanol	nd	N/A
						2-Pentylfuran	nd	N/A
						2-Octen-1-ol	nd	N/A
						2-Methoxy-3-1(methylethyl) pyrazine	nd	N/A
						2-Nonanone	nd	N/A
						Fenchone	nd	N/A
						2-Methyl-isoborneol	nd	N/A
						a-Terpineol	nd	N/A
						Borneol	nd	N/A
						Geosmin	nd	N/A
						Thujopsene	nd	N/A

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

mg/M³: Milligrams per cubic meter

nd: Not detected

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

APPENDIX A

TABLE 20803001-143
MICROBIAL VOLATILE ORGANIC COMPOUNDS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 7, 2008

Page 2

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	START/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (mg/M³)	PEL (mg/M³)
Area Sample	Column N22 area; Cubicle 37; about center;	N/A	20804001-	14:02/	94	3-Methylfuran	nd	N/A
'	approximately six feet above floor/Normal office		M02JL	15:36	minutes	2-Methyl-1-propanol	nd	N/A
	activities					1-Butanol	530 x10 ⁻⁶	300
						3-Methyl-2-butanol	nd	N/A
						2-Pentanol	nd	N/A
						3-Methyl-2-butanol	nd	N/A
						Methyl disulfide	nd	N/A
						Ethyl isobutyrate	nd	N/A
						2-Hexanone	nd	410
						2-Heptanone	nd	468
						5-Methyl-3-heptanone	nd	N/A
						1-Octen-3-ol	nd	N/A
						3-Octanone	nd	N/A
						3-Octanol	nd	N/A
						2-Pentylfuran	nd	N/A
						2-Octen-1-ol	nd	N/A
						2-Methoxy-3-1(methylethyl) pyrazine	nd	N/A
						2-Nonanone	nd	N/A
						Fenchone	nd	N/A
						2-Methyl-isoborneol	nd	N/A
						a-Terpineol	nd	N/A
						Borneol	nd	N/A
						Geosmin	nd	N/A
						Thujopsene	nd	N/A

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

mg/M³: Milligrams per cubic meter

nd: Not detected

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

APPENDIX A

TABLE 20803001-143
MICROBIAL VOLATILE ORGANIC COMPOUNDS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 7, 2008

Page 3

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	START/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (mg/M³)	PEL (mg/M³)
Area Sample	Area between Columns N18 and N19; Cubicle	N/A	20804001-	14:04/	96	3-Methylfuran	nd	N/A
'	122; about center; approximately six feet above		M03JL	15:40	minutes	2-Methyl-1-propanol	nd	N/A
	floor/Normal office activities					1-Butanol	nd	300
						3-Methyl-2-butanol	nd	N/A
						2-Pentanol	nd	N/A
						3-Methyl-2-butanol	nd	N/A
						Methyl disulfide	nd	N/A
						Ethyl isobutyrate	nd	N/A
						2-Hexanone	nd	410
						2-Heptanone	nd	468
						5-Methyl-3-heptanone	nd	N/A
						1-Octen-3-ol	nd	N/A
						3-Octanone	nd	N/A
						3-Octanol	nd	N/A
						2-Pentylfuran	nd	N/A
						2-Octen-1-ol	nd	N/A
						2-Methoxy-3-1(methylethyl) pyrazine	nd	N/A
						2-Nonanone	nd	N/A
						Fenchone	nd	N/A
						2-Methyl-isoborneol	nd	N/A
						a-Terpineol	nd	N/A
						Borneol	nd	N/A
						Geosmin	nd	N/A
						Thujopsene	nd	N/A

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

mg/M³: Milligrams per cubic meter

nd: Not detected

HYGIENE TECHNOLOGIES INTERNATIONAL, INC.

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279

APPENDIX A

TABLE 20803001-143
MICROBIAL VOLATILE ORGANIC COMPOUNDS

2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 7, 2008

Page 4

NAME/ REFERENCE	LOCATION/ ACTIVITIES	PPE USED	SAMPLE NUMBER	START/ STOP	SAMPLE TIME	CONTAMINANT	RESULTS (mg/M³)	PEL (mg/M³)
Area Sample	Area between Columns K19 and K20; about two	N/A	20804001-	14:10	93	3-Methylfuran	nd	N/A
	feet east of Cubicle 20; approximately six feet		M04JL	15:43	minutes	2-Methyl-1-propanol	nd	N/A
	above floor/Normal office activities					1-Butanol	525 x10 ⁻⁶	300
						3-Methyl-2-butanol	nd	N/A
						2-Pentanol	nd	N/A
						3-Methyl-2-butanol	nd	N/A
						Methyl disulfide	nd	N/A
						Ethyl isobutyrate	nd	N/A
						2-Hexanone	nd	410
						2-Heptanone	nd	468
						5-Methyl-3-heptanone	nd	N/A
						1-Octen-3-ol	nd	N/A
						3-Octanone	nd	N/A
						3-Octanol	nd	N/A
						2-Pentylfuran	nd	N/A
						2-Octen-1-ol	nd	N/A
						2-Methoxy-3-1(methylethyl) pyrazine	nd	N/A
						2-Nonanone	nd	N/A
						Fenchone	nd	N/A
						2-Methyl-isoborneol	nd	N/A
						a-Terpineol	nd	N/A
						Borneol	nd	N/A
						Geosmin	nd	N/A
						Thujopsene	nd	N/A

LEGEND

PPE: Personal protective equipment

N/A: Not applicable

PEL: Cal-OSHA 8-hour time-weighted average permissible exposure limit

mg/M³: Milligrams per cubic meter

nd: Not detected

HYGIENE TECHNOLOGIES INTERNATIONAL, INC.

CLIENT: California State Board of Equalization 450 N Street Sacramento, California 94279



TABLE 20803001-144
DIRECT-READING RESULTS
2ND FLOOR
SACRAMENTO, CALIFORNIA
APRIL 8, 2008

LOCATION/SITE ACTIVITIES	SAMPLE TIME	CONTAMINANT	RESULTS (ppm)	COMMENTS
Column K22 area; approximately five feet above floor/Normal office activities	10:30/10:35	Volatile Organic Compounds	0.3	N/A
		Ozone	ND < 0.05	
Column K18 area; about five feet above floor/Normal office activities	10:41/10:47	Volatile Organic Compounds	0.3	N/A
		Ozone	ND < 0.05	
Column N18 area; approximately five feet above floor/Normal office activities	10:52/10:56	Volatile Organic Compounds	0.3	N/A
		Ozone	ND < 0.05	
Column N22 area; approximately five feet above floor/Normal office activities	11:01/11:07	Volatile Organic Compounds	0.3	N/A
		Ozone	ND < 0.05	

ND: Not detected <: Less than

N/A: Not applicable ppm: Parts per million



Report for:

Mr. Wes Frey Hygiene Technologies International, Inc.: Northern California 3127 Bowen Island Street West Sacramento, CA 95691

Regarding: Project: 20804001

EMĹ ID: 410940

Approved by:

h Managar

Lab Manager Magzoub Ismail Dates of Analysis:

Spore trap analysis: 04-17-2008

Project SOPs: Spore trap analysis (I100000)

This coversheet is included with your report in order to comply with AIHA and ISO accreditation requirements.

For clarity, we report the number of significant digits as calculated; but, due to the nature of this type of biological data, the number of significant digits that is used for interpretation should generally be one or two. All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank corrections of results is not a standard practice. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Northern California Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		20804001- TM17outJL		20804001-TM18JL		01-TM19JL	2080400	01-TM20JL
Comments (see below)	N	Vone	N	Vone	N	Vone	N	Vone
Lab ID-Version‡:	180	5509-1	180	5510-1	1805511-1		180	5512-1
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria	2	27		_		_		
Arthrinium								
Ascospores*	1	53						
Aureobasidium								
Basidiospores*	6	320						
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium	9	480						
Curvularia								
Epicoccum								
Fusarium								
Myrothecium								
Nigrospora								
Oidium	1	13						
Other colorless								
Penicillium/Aspergillus types†	8	427	1	53	1	53	1	53
Pithomyces								
Rusts*	1	13						
Smuts*, Periconia, Myxomycetes*	9	120						
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	3+		1+		1+		1+	
Hyphal fragments/m3	360		< 13		< 13		< 13	
Pollen/m3	107		< 13		< 13		13	
Skin cells (1-4+)	1+		2+		1+		2+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		1,453		53		53		53

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens. † The spores of *Aspergillus* and *Penicillium* (and others such as *Acremonium*, *Paecilomyces*) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Date of Sampling: 04-14-2008 Client: Hygiene Technologies International, Inc.: Northern California Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	2080400	1-TM21JL	2080400	1-TM22JL	2080400	1-TM23JL	2080400)1-TM24JL
Comments (see below)	N	lone	N	lone	N	Vone	N	Vone
Lab ID-Version‡:	180.	5513-1	180	5514-1	1805515-1		180	5516-1
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria								
Arthrinium								
Ascospores*								
Aureobasidium								
Basidiospores*							1	53
Bipolaris/Drechslera group								
Botrytis								
Chaetomium					1	13		
Cladosporium			1	53			1	53
Curvularia								
Epicoccum								
Fusarium								
Myrothecium								
Nigrospora								
Oidium							1	13
Other colorless								
Penicillium/Aspergillus types†					1	53		
Pithomyces								
Rusts*								
Smuts*, Periconia, Myxomycetes*	2	27						
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	1+		1+		1+		2+	
Hyphal fragments/m3	< 13		< 13		< 13		< 13	
Pollen/m3	13		13		27		27	
Skin cells (1-4+)	1+		2+		2+		2+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		27		53		66		119

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

^{*}Most of these spore types are not seen with culturable methods (Andersen sampling), annough some may appear as not sportating rengt.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

† The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Hygiene Technologies International, Inc.:

Northern California

C/O: Mr. Wes Frey

Date of Sampling: 04-14-2008

Date of Receipt: 04-15-2008

Date of Report: 04-17-2008

Re: 20804001

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	2080400	1-TM25JL	2080400	1-TM26JL	2080400	1-TM27JL	2080400)1-TM28JL
Comments (see below)	N	lone	N	Ione		A		В
Lab ID-Version‡:	180	5517-1	180	5518-1	180	5519-1	180	5520-1
	raw ct.	spores/m3						
Alternaria								
Arthrinium								
Ascospores*								
Aureobasidium								
Basidiospores*			1	53	1	53		
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium	1	53			1	53	1	53
Curvularia								
Epicoccum								
Fusarium								
Myrothecium								
Nigrospora								
Oidium								
Other colorless								
Penicillium/Aspergillus types†	2	107	1	53	36	880	10	293
Pithomyces								
Rusts*					1	13		
Smuts*, Periconia, Myxomycetes*					5	67		
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	2+		2+		2+		2+	
Hyphal fragments/m3	< 13		< 13		< 13		< 13	
Pollen/m3	13		< 13		53		53	
Skin cells (1-4+)	2+		2+		2+		2+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		160		106		1,066		346

Comments: A) 26 of the raw count *Penicillium/Aspergillus* type spores were present as a single clump. B) 6 of the raw count *Penicillium/Aspergillus* type spores were present as a single clump.

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Hygiene Technologies International, Inc.:
Northern California
C/O: Mr. Wes Frey

Date of Sampling: 04-14-2008
Date of Receipt: 04-15-2008
Date of Report: 04-17-2008

Re: 20804001

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	2080400	1-TM29JL	2080400	1-TM30JL	2080400	1-TM31JL	20804001-TM32J	
Comments (see below)	N	Ione	N	Ione		С	N	Vone
Lab ID-Version‡:	180	5521-1	180:	5522-1	180	5523-1	180	5524-1
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria								
Arthrinium								
Ascospores*								
Aureobasidium								
Basidiospores*							1	53
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium	2	107	3	160				
Curvularia								
Epicoccum								
Fusarium								
Myrothecium								
Nigrospora								
Oidium								
Other colorless								
Penicillium/Aspergillus types†	1	53					1	53
Pithomyces								
Rusts*								
Smuts*, Periconia, Myxomycetes*			2	27			1	13
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	1+		1+		1+		2+	
Hyphal fragments/m3	< 13		< 13		< 13		13	
Pollen/m3	< 13		27		13		40	
Skin cells (1-4+)	2+		2+		1+		2+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		160		187		< 13		119

Comments: C) No spores detected.

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Date of Sampling: 04-14-2008 Client: Hygiene Technologies International, Inc.: Northern California Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	208040	01-TM33JL	20804001	-TM34outJL	
Comments (see below)]	None	N	lone	
Lab ID-Version‡:	180)5525-1	1805526-1		
	raw ct.	spores/m3	raw ct.	spores/m3	
Alternaria			1	13	
Arthrinium					
Ascospores*					
Aureobasidium					
Basidiospores*	3	160	5	267	
Bipolaris/Drechslera group					
Botrytis					
Chaetomium					
Cladosporium	1	53	40	2,130	
Curvularia			1	13	
Epicoccum					
Fusarium					
Myrothecium					
Nigrospora					
Oidium			1	13	
Other colorless					
Penicillium/Aspergillus types†	1	53	14	747	
Pithomyces					
Rusts*			1	13	
Smuts*, Periconia, Myxomycetes*			15	200	
Stachybotrys			2	27	
Stemphylium					
Torula			3	40	
Ulocladium					
Zygomycetes					
Background debris (1-4+)††	1+		3+		
Hyphal fragments/m3	< 13		120		
Pollen/m3	13		27		
Skin cells (1-4+)	1+		1+		
Sample volume (liters)	75		75		
TOTAL SPORE/m3		266		3,463	

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens.

† The spores of *Aspergillus* and *Penicillium* (and others such as *Acremonium*, *Paecilomyces*) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: 20804001-TM17outJL

Fungi Identified	Outdoor	Туріса	al Outdoo	r Data by	Outdoor	Data by L	ocation‡		
	data		Month	ı: April		State: CA			
	spores/m3	low	med	high	freq %	low	med	high	freq %
Generally able to grow indoors*									
Alternaria	27	7	27	230	49	7	27	230	60
Bipolaris/Drechslera group	-	7	13	130	13	7	13	120	14
Chaetomium	-	7	13	130	13	7	13	110	19
Cladosporium	480	27	370	4,200	93	53	640	6,500	98
Curvularia	-	7	13	210	7	7	13	210	7
Nigrospora	-	7	13	93	7	7	13	170	8
Penicillium/Aspergillus types	427	27	160	1,500	79	40	210	2,500	88
Stachybotrys	-	7	13	480	3	7	13	310	5
Torula	-	7	13	160	12	7	13	150	13
Seldom found growing indoors**									
Ascospores	53	13	110	2,500	75	13	110	1,800	73
Basidiospores	320	13	210	5,200	91	13	250	6,800	95
Oidium	13	7	20	230	22	7	13	190	20
Rusts	13	7	20	250	24	7	13	270	29
Smuts, Periconia, Myxomycetes	120	7	33	410	62	8	40	470	71
TOTAL SPORES/M3	1,453								

[†] The Typical Outdoor Data by Date represents the typical outdoor spore levels across North America for the month indicated. The last column represents the frequency of occurrence. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 2.5% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

[‡] The Typical Outdoor Data by Location represents the typical outdoor spore levels for the region indicated for the entire year. As with the Typical Outdoor Data by Date, the four columns represent the frequency of occurrence and the typical low, medium, and high concentration values for the spore type indicated. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

^{*}The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**}These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Re: 20804001 Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldRANGETM: Extended Outdoor Comparison

Outdoor Location: 20804001-TM34outJL

Fungi Identified	Outdoor	Туріса	al Outdoo	r Data by	Date†	Typical Outdoor Data by Location:			
	data		Month	: April		State: CA			
	spores/m3	low	med	high	freq %	low	med	high	freq %
Generally able to grow indoors*									
Alternaria	13	7	27	230	49	7	27	230	60
Bipolaris/Drechslera group	-	7	13	130	13	7	13	120	14
Chaetomium	-	7	13	130	13	7	13	110	19
Cladosporium	2,130	27	370	4,200	93	53	640	6,500	98
Curvularia	13	7	13	210	7	7	13	210	7
Nigrospora	-	7	13	93	7	7	13	170	8
Penicillium/Aspergillus types	747	27	160	1,500	79	40	210	2,500	88
Stachybotrys	27	7	13	480	3	7	13	310	5
Torula	40	7	13	160	12	7	13	150	13
Seldom found growing indoors**									
Ascospores	-	13	110	2,500	75	13	110	1,800	73
Basidiospores	267	13	210	5,200	91	13	250	6,800	95
Oidium	13	7	20	230	22	7	13	190	20
Rusts	13	7	20	250	24	7	13	270	29
Smuts, Periconia, Myxomycetes	200	7	33	410	62	8	40	470	71
TOTAL SPORES/M3	3,463								

[†] The Typical Outdoor Data by Date represents the typical outdoor spore levels across North America for the month indicated. The last column represents the frequency of occurrence. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values of the spore type when it is detected. For example, if the frequency of occurrence is 63% and the low value is 53, it would mean that the given spore type is detected 63% of the time and, when detected, 2.5% of the time it is present in levels above the detection limit and below 53 spores/m3. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor data" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. In addition, EMLab P&K may not have received and tested a representative number of samples for every region or time period. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the use or interpretation of the data contained in, or any actions taken or omitted in reliance upon, this report.

[‡] The Typical Outdoor Data by Location represents the typical outdoor spore levels for the region indicated for the entire year. As with the Typical Outdoor Data by Date, the four columns represent the frequency of occurrence and the typical low, medium, and high concentration values for the spore type indicated. These values are updated periodically, and if enough data is not available to make a statistically meaningful assessment, it is indicated with a dash.

^{*}The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**}These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Northern California C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 20804001-TM17outJL:

Species detected		Outdoor	r sample sp	ores/m3		Typical	outdo	or ranges	Freq.
	<100	1K	10K	>100K		(North America)			%
Alternaria					27	7 -	27	- 380	54
Ascospores					53	13 -	150	- 4,200	75
Basidiospores					320	13 -	320	- 14,000	92
Cladosporium					480	38 -	530	- 8,400	94
Oidium					13	7 -	13	- 220	15
Penicillium/Aspergillus types					427	27 -	210	- 2,500	85
Rusts					13	7 -	13	- 310	23
Smuts, Periconia, Myxomycetes					120	7 -	40	- 750	70
Total					1,453				

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 20804001-TM18JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ment ratio** or/outdoor)	Spearman ran correlation*** (indoor/outdoo	(indoor/outdoor)
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Re	sult: 0.2222	dF: 8 Result: 0.5774 Critical value: 0.61 Outside Similar: N	
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
Penic	illium/Aspergillus types				53
	Total				53

Client: Hygiene Technologies International, Inc.: Northern California

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM19JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		correl	nan rank ation*** /outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.2222		Resul Critical v	F: 8 t: 0.5774 alue: 0.6190 Similar: No	Score: 107 Result: Low
Species 1	Detected			Spor	res/m3	
		<100	1K		10K	>100K
Penic	illium/Aspergillus types					53
	Total					53

Location: 20804001-TM20JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.2222		dF: 8 Result: 0.5774 Critical value: 0.6190 Outside Similar: No		Score: 107 Result: Low
Species	Detected			Spore	es/m3	
		<100	1K		10K	>100K
Penic	illium/Aspergillus types					53
	Total					53

Location: 20804001-TM21JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ement ratio** oor/outdoor)	correl	nan rank ation*** ⁄outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Re	esult: 0.2222	Resul Critical v	IF: 8 t: 0.3869 value: 0.6190 Similar: No	Score: 105 Result: Low
Species	Detected			Spo	res/m3	
		<100	1K		10K	>100K
Smuts, F	Periconia, Myxomycetes					27
	Total					27

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Re: 20804001 Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM22JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE*** (indoor/outdoor)
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.2222		dF: 8 Result: 0.6726 Critical value: 0.6190 Outside Similar: Yes		Score: 103 Result: Low
Species 1	Detected			Spore	s/m3	
		<100	1K		10K	>100K
	Cladosporium					53
	Total					53

Location: 20804001-TM23JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)
Result: 4%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.2000		dF: 9 Result: 0.2208 Critical value: 0.5833 Outside Similar: No		Score: 121 Result: Low
Species	Detected			Spores/		
		<100	1K		10K	>100K
	Chaetomium					13
Penicillium/Aspergillus types						53
	Total					66

Location: 20804001-TM24JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 8%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.5455	dF: 8 Result: 0.4524 Critical value: 0.6190 Outside Similar: No	Score: 104 Result: Low
Species 1	Detected		Spores/m3	
		<100 1K	10K	>100K
	Basidiospores			53
	Cladosporium			53
	Oidium			13
	Total			119

Client: Hygiene Technologies International, Inc.: Northern California Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM25JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORI (indoor/outo	
Result: 11%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4000		dF: 8 Result: 0.7738 Critical value: 0.6190 Outside Similar: Yes		Score: 112 Result: Lo	_
Species 1	Detected			Spores	s/m3		
		<100	1K		10K	>100K	
	Cladosporium						53
Penic	illium/Aspergillus types						107
	Total						160

Location: 20804001-TM26JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 7%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4000		dF: 8 Result: 0.6012 Critical value: 0.6190 Outside Similar: No	Score: 105 Result: Low
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
	Basidiospores				53
Penicillium/Aspergillus types					53
	Total				106

Client: Hygiene Technologies International, Inc.: Northern California

C/O: Mr. Wes Frey

Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Date of Sampling: 04-14-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM27JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)		
Result: 73%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.7692		dF: 8 Result: 0.7262 Critical value: 0.6190 Outside Similar: Yes	Score: 197 Result: Medium		
Species Detected		Spores/m3					
		<100 1K		10K	>100K		
	Basidiospores				53		
	Cladosporium				53		
Penici	illium/Aspergillus types				880		
Rusts					13		
Smuts, Periconia, Myxomycetes					67		
,	Total				1,066		

Location: 20804001-TM28JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ement ratio** oor/outdoor)	corre	man rank elation*** or/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 23%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Re	esult: 0.4000	Resu Critical	dF: 8 alt: 0.7738 value: 0.6190 e Similar: Yes	Score: 135 Result: Low
Species	Detected			Spo	ores/m3	
_		<100	1K	_	10K	>100K
	Cladosporium					53
Penic	illium/Aspergillus types					293
	Total					346

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM29JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORI (indoor/outo	
Result: 11%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4000		dF: 8 Result: 0.7976 Critical value: 0.6190 Outside Similar: Yes		Score: 104 Result: Lo	-
Species 1	Detected			Spor	es/m3		
		<100	1K		10K	>100K	
	Cladosporium						107
Penic	illium/Aspergillus types						53
	Total						160

Location: 20804001-TM30JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 12%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4000	dF: 8 Result: 0.6310 Critical value: 0.6190 Outside Similar: Yes	Score: 107 Result: Low
Species 1	Detected		Spores/m3	
		<100 1K	10K	>100K
	Cladosporium			160
Smuts, P	Periconia, Myxomycetes			27
	Total			187

Location: 20804001-TM31JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement (indoor/or		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: < 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.0000		dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low
Species 1	Detected			Spores/m3	
		<100	1K	10K	>100K
	None Detected				N/A

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Northern California Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM32JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 8%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.5455		dF: 8 Result: 0.5952 Critical value: 0.6190 Outside Similar: No	Score: 104 Result: Low	
Species 1	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Basidiospores				53	
Penicillium/Aspergillus types					53	
Smuts, Periconia, Myxomycetes					13	
	Total				119	

Location: 20804001-TM33JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ra (indoor/outd		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)
Result: 18%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.545	55	dF: 8 Result: 0.8095 Critical value: 0.6190 Outside Similar: Yes	Score: 112 Result: Low
Species 1	Detected			Spores/m3	
		<100	1K	10K	>100K
	Basidiospores				160
	Cladosporium				53
Penicillium/Aspergillus types					53
	Total				266

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Date of Sampling: 04-14-2008

Date of Receipt: 04-15-2008

Date of Report: 04-17-2008

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab ID: 410940, Page 8 of 8

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Outdoor Summary: 20804001-TM34outJL:

Species detected		Outdoor	sample s	pores/m3	Typical	Typical outdoor ranges		
	<100	1K	10K	>100K	(No	(North America)		
Alternaria				13	7 -	27	- 380	54
Ascospores				ND	13 -	150	- 4,200	75
Basidiospores				267	13 -	320	- 14,000	92
Cladosporium				2,130	38 -	530	- 8,400	94
Curvularia				13] 7 -	22	- 640	16
Oidium				13] 7 -	13	- 220	15
Penicillium/Aspergillus types				747] 27 -	210	- 2,500	85
Rusts				13] 7 -	13	- 310	23
Smuts, Periconia, Myxomycetes				200] 7 -	40	- 750	70
Stachybotrys				27] 7 -	13	- 350	3
Torula				40] 7 -	13	- 160	12
Total				3,463				

The "Typical outdoor ranges" and "Freq. %" columns show the typical low, medium, and high spore counts per cubic meter and the frequency of occurrence for the given spore type. The low, medium, and high values represent the 2.5, 50, and 97.5 percentile values when the spore type is detected. For example, if the low value is 53 and the frequency of occurrence is 63%, it would mean that we typically detect the given spore type on 63 percent of all outdoor samples and, when detected, 2.5% of the time it is present in levels below 53 spores/m3.

Indoor Samples

Location: 20804001-TM18JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Res	sult: 0.1818	Result Critical va	7: 10 : 0.6061 due: 0.5515 imilar: Yes	Score: 107 Result: Low	
Species	Detected			Spor	es/m3		
		<100	1K		10K	>100K	
Penic	illium/Aspergillus types					5	53
	Total					5	53

Client: Hygiene Technologies International, Inc.: Northern California

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM19JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.1818		dF: 10 Result: 0.6061 Critical value: 0.5515 Outside Similar: Yes		Score: 107 Result: Low	
Species 1	Detected	Spores/m3					
		<100	1K		10K	>100K	
Penic	illium/Aspergillus types					53	
	Total					53	

Location: 20804001-TM20JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Res	ult: 0.1818	dF: Result: (Critical valu Outside Sin	0.6061 ne: 0.5515	Score: 107 Result: Low
Species 1	Detected	Spores/m3				
		<100	1K		10K	>100K
Penic	illium/Aspergillus types					53
	Total					53

Location: 20804001-TM21JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: < 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Re	esult: 0.1818	Resu Critical	dF: 10 dt: 0.4848 value: 0.5515 e Similar: No	Score: 105 Result: Low	
Species	Detected			Spo	ores/m3		
		<100	1K		10K	>100K	
Smuts, F	Periconia, Myxomycetes					27	
	Total					27	

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Re: 20804001 Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM22JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor	(indoor/outdoor)
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.1818		dF: 10 Result: 0.6667 Critical value: 0.551 Outside Similar: Ye	-
Species	Detected			Spores/m3	
		<100	1K	10K	>100K
	Cladosporium				53
	Total				53

Location: 20804001-TM23JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.1667		dF: 11 Result: 0.2864 Critical value: 0.5273 Outside Similar: No		Score: 121 Result: Low	
Species 1	Detected			Spor	res/m3		
		<100	1K		10K	>100K	
	Chaetomium					13	
Penicillium/Aspergillus types						53	
	Total					66	

Location: 20804001-TM24JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4615		dF: 10 Result: 0.5242 Critical value: 0.5515 Outside Similar: No	Score: 105 Result: Low	
Species 1	Detected			Spores/m3		
		<100	1K	10K	>100K	
	Basidiospores				53	
	Cladosporium				53	
	Oidium				13	
	Total				119	

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Northern California C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM25JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)		Spearman rank correlation*** (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)	
Result: 4%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.33	33	Result Critical va	F: 10 t: 0.7636 alue: 0.5515 Similar: Yes	Score: 11 Result: Lo	
Species 1	Detected			Spor	res/m3		
		<100	1K		10K	>100K	
	Cladosporium						53
Penic	llium/Aspergillus types						107
	Total						160

Location: 20804001-TM26JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio** (indoor/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)	
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.6515 Critical value: 0.5515 Outside Similar: Yes	Score: 105 Result: Low	
Species 1	Detected		Spores/m3		
		<100 1K	10K	>100K	
	Basidiospores			53	
Penicillium/Aspergillus types				53	
	Total			106	

Client: Hygiene Technologies International, Inc.: Northern California Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM27JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ra (indoor/outd		Spearman correlatio (indoor/ou	n***	MoldSCORE**** (indoor/outdoor)				
Result: 30%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.666	57	dF: 10 Result: 0. Critical value Outside Simi	7727 : 0.5515	Score: 197 Result: Medium				
Species 1	Spores/m3									
		<100	1K		10K	>100K				
	Basidiospores						53			
	Cladosporium						53			
Penici	illium/Aspergillus types						880			
Rusts							13			
Smuts, Periconia, Myxomycetes							67			
,	Total						1,066			

Location: 20804001-TM28JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ement ratio** oor/outdoor)	correl	nan rank lation*** r/outdoor)	MoldSCORE**** (indoor/outdoor)			
Result: 9%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Re	esult: 0.3333	Resul Critical v	F: 10 lt: 0.7636 value: 0.5515 Similar: Yes	Score: 135 Result: Low			
Species	Detected			Spo	res/m3				
_		<100	1K		10K	>100K			
	Cladosporium					53			
Penic	illium/Aspergillus types					293			
	Total					346			

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Northern California C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM29JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio (indoor/outdoor		MoldSCORE**** (indoor/outdoor)
Result: 4%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.7758 Critical value: 0.5515 Outside Similar: Yes	
Species 1	Detected		Spores/m3	
		<100	1K 10K	>100K
	Cladosporium			107
Penicillium/Aspergillus types				53
	Total			160

Location: 20804001-TM30JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio (indoor/outdoor		MoldSCORE**** (indoor/outdoor)		
Result: 5%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.3333	dF: 10 Result: 0.6667 Critical value: 0.5515 Outside Similar: Yes	Score: 103 Result: Low		
Species 1	Detected		Spores/m3			
		<100	K 10K	>100K		
	Cladosporium			160		
Smuts, Periconia, Myxomycetes				27		
	Total			187		

Location: 20804001-TM31JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		ment ratio** or/outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)					
Result: < 1%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Res	sult: 0.0000	dF: N/A Result: N/A Critical value: N/A Outside Similar: N/A	Score: 100 Result: Low					
Species	Detected	Spores/m3								
		<100	1K	10K	>100K					
	None Detected				N/A					

Client: Hygiene Technologies International, Inc.:

Date of Sampling: 04-14-2008 Northern California Date of Receipt: 04-15-2008 C/O: Mr. Wes Frey Date of Report: 04-17-2008

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

Location: 20804001-TM32JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)		nt ratio** outdoor)	Spearman rank correlation*** (indoor/outdoor)	MoldSCORE**** (indoor/outdoor)				
Result: 3%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result	: 0.4615	dF: 10 Result: 0.6758 Critical value: 0.5515 Outside Similar: Yes	Score: 105 Result: Low				
Species 1	Detected	Spores/m3							
		<100	1K	10K	>100K				
	Basidiospores				53				
Penicillium/Aspergillus types					53				
Smuts, Periconia, Myxomycetes					13				
	Total				119				

Location: 20804001-TM33JL

% of outdoor total spores/m3	Friedman chi- square* (indoor variation)	Agreement ratio* (indoor/outdoor)		MoldSCORE**** (indoor/outdoor)					
Result: 7%	dF: 15 Result: 10.4622 Critical value: 24.9958 Inside Similar: Yes	Result: 0.4615	dF: 10 Result: 0.8212 Critical value: 0.5515 Outside Similar: Yes	Score: 115 Result: Low					
Species 1	Detected	Spores/m3							
		<100	K 10K	>100K					
	Basidiospores			160					
Cladosporium				53					
Penicillium/Aspergillus types				53					
	Total			266					

^{*} The Friedman chi-square statistic is a non-parametric test that examines variation in a set of data (in this case, all indoor spore counts). The null hypothesis (H0) being tested is that there is no meaningful difference in the data for all indoor locations. The alternative hypothesis (used if the test disproves the null hypothesis) is that there is a difference between the indoor locations. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

^{**} An agreement ratio is a simple method for assessing the similarity of two samples (in this case the indoor sample and the outdoor summary) based on the spore types present. A score of one indicates that the types detected in one location are the same as that in the other. A score of zero indicates that none of the types detected indoors are present outdoors. Typically, an agreement of 0.8 or higher is considered high.

^{***} The Spearman rank correlation is a non-parametric test that examines correlation between two sets of data (in this case the indoor location and the outdoor summary). The null hypothesis (H0) being tested is that the indoor and outdoor samples are unrelated. The alternative hypothesis (used if the test disproves the null hypothesis) is that the samples are similar. The null hypothesis is rejected when the result of the test is greater than the critical value. The critical value that is displayed is based on the degrees of freedom (dF) of the test and a significance level of 0.05.

Date of Sampling: 04-14-2008

Date of Receipt: 04-15-2008

Date of Report: 04-17-2008

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

MoldSTATTM: Supplementary Statistical Spore Trap Report

**** MoldSCORETM is a specialized method for examining air sampling data. It is a score between 100 and 300, with 100 indicating a greater likelihood that the airborne indoor spores originated from the outside, and 300 indicating a greater likelihood that they originated from an inside source. The Result displayed is based on the numeric score given and will be either Low, Medium, or High, indicating a low, medium, or high likelihood that the spores detected originated from an indoor source. EMLab P&Kreserves the right to, and may at anytime, modify or change the MoldScore algorithm without notice.

Interpretation of the data contained in this report is left to the client or the persons who conducted the field work. This report is provided for informational and comparative purposes only and should not be relied upon for any other purpose. "Typical outdoor ranges" are based on the results of the analysis of samples delivered to and analyzed by EMLab P&K and assumptions regarding the origins of those samples. Sampling techniques, contaminants infecting samples, unrepresentative samples and other similar or dissimilar factors may affect these results. With the statistical analysis provided, as with all statistical comparisons and analyses, false-positive and false-negative results can and do occur. EMLab P&K hereby disclaims any liability for any and all direct, indirect, punitive, incidental, special or consequential damages arising out of the data contained in, or any actions taken or omitted in reliance upon, this report.

EMLab ID: 410940, Page 8 of 8

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report Outdoor Sample: 20804001-TM17outJL

Fungi Identified	Οι	ıto	do	or	san	ıpl	le :	spo	res	/m	3	Raw	Spores/
Ü	<10	0		1	K			10K		>100	0 K	count	m3
Generally able to grow indoors*													
Alternaria												2	27
Bipolaris/Drechslera group												ND	< 13
Chaetomium							Ш					ND	< 13
Cladosporium												9	480
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												8	427
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores††												1	53
Basidiospores††												6	320
Oidium												1	13
Rusts												1	13
Smuts, Periconia, Myxomycetes††												9	120
Total													1,453

Location: 20804001-TM18JL

Fungi Identified	In	do	or	sam	pl	e s	spo	res	/n	13	Raw	Spores/
	<100)		1K			10 F	ζ	>1	00k	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											ND	< 13
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											1	53
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores††											ND	< 13
Basidiospores††											ND	< 13
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												53

100	MoldSCORE 100 200 300									
100	200	300	Score							
			100							
			100							
			100							
			100							
			100							
			100							
			107							
			100							
			100							
			100							
			100							
			100							
			100							
Fina	l MoldSC(ORE	107							

Client: Hygiene Technologies International, Inc.: Northern California

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM19JL

Fungi Identified	Inde	00	r	samj	ple	sp	ore	es/1	m.	3	Raw	Spores/
	<100			1K		1	0K	>	>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											ND	< 13
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											1	53
Stachybotrys											ND	< 13
Torula		П									ND	< 13
Seldom found growing indoors**												
Ascospores††		П							Ш		ND	< 13
Basidiospores††		П									ND	< 13
Rusts		П									ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												53

100 MoldSCORE :	‡ Score
	100
	100
	100
	100
	100
	100
	107
	100
	100
	100
	100
	100
	100
Final MoldSCORE	107

Location: 20804001-TM20JL

Fungi Identified	Indo	or sam	ple spore	s/m3	Raw	Spores/
	<100	1K	10K	>100K	count	m3
Generally able to grow indoors*						
Alternaria					ND	< 13
Bipolaris/Drechslera group					ND	< 13
Chaetomium					ND	< 13
Cladosporium					ND	< 13
Curvularia					ND	< 13
Nigrospora					ND	< 13
Penicillium/Aspergillus types†					1	53
Stachybotrys					ND	< 13
Torula					ND	< 13
Seldom found growing indoors**						
Ascospores††					ND	< 13
Basidiospores††					ND	< 13
Rusts					ND	< 13
Smuts, Periconia, Myxomycetes††					ND	< 13
Total						53

100	Score								
	100 200 300								
			100						
			100						
			100						
			100						
			100						
			100						
			107						
			100						
			100						
			100						
			100						
			100						
			100						
Fina	al MoldSC	ORE	107						

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM21JL

Fungi Identified	In	do	001	r s	sam	ple	e s	por	es/	m.	3	Raw	Spores/
	<10	0			1K			10K		>10	0K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group												ND	< 13
Chaetomium												ND	< 13
Cladosporium												ND	< 13
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												ND	< 13
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores††												ND	< 13
Basidiospores††												ND	< 13
Rusts												ND	< 13
Smuts, Periconia, Myxomycetes††							П					2	27
Total													27

MoldS (Score
		100
		100
		100
		100
		100
		100
		100
		100
		100
		100
		100
		100
		105
Final MoldS	CORE	105

Location: 20804001-TM22JL

Fungi Identified	Ind	oor sa	mple	spore	es/m3	Raw	Spores/
	<100	11	ζ	10K	>100K	count	m3
Generally able to grow indoors*							
Alternaria						ND	< 13
Bipolaris/Drechslera group						ND	< 13
Chaetomium						ND	< 13
Cladosporium						1	53
Curvularia						ND	< 13
Nigrospora						ND	< 13
Penicillium/Aspergillus types†						ND	< 13
Stachybotrys						ND	< 13
Torula						ND	< 13
Seldom found growing indoors**							
Ascospores††						ND	< 13
Basidiospores††						ND	< 13
Rusts						ND	< 13
Smuts, Periconia, Myxomycetes††						ND	< 13
Total							53

100	Score								
	100 200 300								
			100						
			100						
			100						
			103						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
Fina	Final MoldSCORE								

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Re: 20804001 Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM23JL

Fungi Identified	Indo	Indoor sample spores/m3						Raw	Spores/
_	<100		1K		10K	>	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								1	13
Cladosporium								ND	< 13
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total			·						66

MoldSCORE;	Score
	100
	100
	121
	100
	100
	100
	106
	100
	100
	100
	100
	100
	100
Final MoldSCORE	121

Location: 20804001-TM24JL

Fungi Identified	Inc	door	r sa	mp	le	spo	res	/m.	3	Raw	Spores/
	<100		1K			10 F	(>10	0K	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										1	53
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores††										ND	< 13
Basidiospores††										1	53
Oidium										1	13
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes††										ND	< 13
Total											119

100	Score					
			100			
			100			
			100			
			102			
			100			
			100			
			100			
			100			
			100			
			100			
			104			
			105			
			100			
			100			
Fina	Final MoldSCORE					

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

C/O: Mr. Wes Frey Dat Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM25JL

Fungi Identified	Indo	Indoor sample spores/m3						Raw	Spores/
	<100		1K		10K	>	100I	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								1	53
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								2	107
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total			•		·				160

MoldSCORE‡								
100 200 300	Score							
	100							
	100							
	100							
	101							
	100							
	100							
	112							
	100							
	100							
	100							
	100							
	100							
	100							
Final MoldSCORE	112							

Location: 20804001-TM26JL

Fungi Identified	Ind	oor	sam	ple s	spor	es/r	n3	Raw	Spores/
	<100		1K		10K	>	100I	count	m3
Generally able to grow indoors*									
Alternaria							Ш	ND	< 13
Bipolaris/Drechslera group				Ш				ND	< 13
Chaetomium								ND	< 13
Cladosporium							Ш	ND	< 13
Curvularia				Ш				ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys				Ш				ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								1	53
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††				Ш				ND	< 13
Total									106

100	ORE:	Score					
			100				
			100				
			100				
			100				
			100				
			100				
			105				
			100				
			100				
			100				
			104				
			100				
			100				
Fina	Final MoldSCORE						

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM27JI

Fungi Identified	Iı	ıdo	01	rs	samp	ole	S	por	es/i	m3	3	Raw	Spores/
	<10	0			1K			10K		>10	0K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group												ND	< 13
Chaetomium												ND	< 13
Cladosporium												1	53
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												36	880
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores††												ND	< 13
Basidiospores††												1	53
Rusts												1	13
Smuts, Periconia, Myxomycetes††												5	67
Total					•								1,066

MoldSCORE;								
100 200	300 Score							
	100							
	100							
	100							
	100							
	100							
	100							
	197							
	100							
	100							
	100							
	100							
	102							
	101							
Final MoldSCO	RE 197							

Location: 20804001-TM28JL

Fungi Identified	Ind	looi	r s	amp	ole	S	pore	es/i	m3		Raw	Spores/
	<100		1	K			10K		>100	K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium									Ш		ND	< 13
Cladosporium											1	53
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											10	293
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores††											ND	< 13
Basidiospores††											ND	< 13
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												346

100	ORE:	Score	
			100
			100
			100
			100
			100
			100
			135
			100
			100
			100
			100
			100
			100
Fina	135		

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM29II

Fungi Identified	Indo	or	samj	ole s	spore	es/n	13	Raw	Spores/
	<100		1K		10K	>1	001	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								2	107
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††						П		ND	< 13
Basidiospores††						П		ND	< 13
Rusts						П		ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									160

MoldSCORE:	Score
100 200 200	
	100
	100
	100
	104
	100
	100
	103
	100
	100
	100
	100
	100
	100
Final MoldSCORE	104

Location: 20804001-TM30JL

Fungi Identified	Ind	oor s	amp	le s	pore	es/m	13	Raw	Spores/
	<100		١K		10K	>1	00K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								3	160
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								2	27
Total									187

100	ORE:	Score					
			100				
			100				
			100				
			107				
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			100				
			103				
Fina	Final MoldSCORE						

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM31JL

Fungi Identified	Indo	or	samj	ple s	spore	es/n	13	Raw	Spores/
_	<100		1K		10K	>1	100k	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								ND	< 13
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									N/A

100	MoldSC		Score					
100	200	200 300						
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
			100					
Fin	100							

Location: 20804001-TM32JL

Fungi Identified	Ind	oor	samj	13	Raw	Spores/		
	<100		1K	10K	>	100K	count	m3
Generally able to grow indoors*								
Alternaria							ND	< 13
Bipolaris/Drechslera group							ND	< 13
Chaetomium							ND	< 13
Cladosporium							ND	< 13
Curvularia							ND	< 13
Nigrospora							ND	< 13
Penicillium/Aspergillus types†							1	53
Stachybotrys							ND	< 13
Torula							ND	< 13
Seldom found growing indoors**								
Ascospores††							ND	< 13
Basidiospores††							1	53
Rusts							ND	< 13
Smuts, Periconia, Myxomycetes††							1	13
Total								119

MoldSCORE;								
100	Score							
	100 200 300							
			100					
			100					
			100					
			100					
			100					
			100					
			104					
			100					
			100					
			100					
			104					
			100					
			101					
Fina	104							

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Date of Sampling: 04-14-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM33JL

Fungi Identified	Indoor sample spores/m3								Raw	Spores/		
	<100			1K			10K	>	100	K	count	m3
Generally able to grow indoors*												
Alternaria	Ш										ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											1	53
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											1	53
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores††											ND	< 13
Basidiospores††											3	160
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												266

100	MoldSCORE; Score									
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		112								
		100								
		100								
Fi	nal MoldSCORE	112								
Fi	nal MoldSCORE	112								

^{*}The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**}These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

[†]The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

^{††}Most of these spore types are not seen with culturable methods (Anderson sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores.

[‡]Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSCORETM: **Spore Trap Report Outdoor Sample:** 20804001-TM34outJL

Fungi Identified	Outdoor sample spores/m3								n3	Raw	Spores/	
_	<10	0		1K			1	10K	>1	00K	count	m3
Generally able to grow indoors*												
Alternaria											1	13
Bipolaris/Drechslera group											ND	< 13
Chaetomium										Ш	ND	< 13
Cladosporium											40	2,130
Curvularia											1	13
Nigrospora										Ш	ND	< 13
Penicillium/Aspergillus types†											14	747
Stachybotrys											2	27
Torula											3	40
Seldom found growing indoors**												
Ascospores††											ND	< 13
Basidiospores††											5	267
Oidium											1	13
Rusts											1	13
Smuts, Periconia, Myxomycetes††											15	200
Total												3,463

Location: 20804001-TM18JL

Fungi Identified	Indoor sample spores/m3									Raw	Spores/
	<100	1	1	K		1	0K	>	100F	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										ND	< 13
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										1	53
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores††										ND	< 13
Basidiospores††										ND	< 13
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes††										ND	< 13
Total											53

100	Score						
			100				
			100				
			100				
			100				
			100				
			100				
			107				
			100				
			100				
			100				
			100				
			100				
			100				
Fina	l MoldSCC	RE	107				

Client: Hygiene Technologies International, Inc.: Northern California

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM19JL

Fungi Identified	Inde	Indoor sample spores/m3							3	Raw	Spores/	
	<100			1K		1	0K	>	>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group											ND	< 13
Chaetomium											ND	< 13
Cladosporium											ND	< 13
Curvularia											ND	< 13
Nigrospora											ND	< 13
Penicillium/Aspergillus types†											1	53
Stachybotrys											ND	< 13
Torula		П									ND	< 13
Seldom found growing indoors**												
Ascospores††		П							Ш		ND	< 13
Basidiospores††		П									ND	< 13
Rusts		П									ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												53

MoldSCORE‡ 200 300 Score										
	100									
	100									
	100									
	100									
	100									
	100									
	107									
	100									
	100									
	100									
	100									
	100									
	100									
Final MoldSCORE	107									

Location: 20804001-TM20JL

Fungi Identified	Indo	or sam	ple spore	s/m3	Raw	Spores/
	<100	1K	10K	>100K	count	m3
Generally able to grow indoors*						
Alternaria					ND	< 13
Bipolaris/Drechslera group					ND	< 13
Chaetomium					ND	< 13
Cladosporium					ND	< 13
Curvularia					ND	< 13
Nigrospora					ND	< 13
Penicillium/Aspergillus types†					1	53
Stachybotrys					ND	< 13
Torula					ND	< 13
Seldom found growing indoors**						
Ascospores††					ND	< 13
Basidiospores††					ND	< 13
Rusts					ND	< 13
Smuts, Periconia, Myxomycetes††					ND	< 13
Total						53

100	MoldSCORE; 100 200 300 Score									
			100							
			100							
			100							
			100							
			100							
			100							
			107							
			100							
			100							
			100							
			100							
			100							
			100							
Fina	al MoldSC	ORE	107							

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM21JL

Fungi Identified	Indoor sa	ample spore	es/m3	Raw	Spores/
_	<100 11	K 10K	>100K	count	m3
Generally able to grow indoors*					
Alternaria				ND	< 13
Bipolaris/Drechslera group				ND	< 13
Chaetomium				ND	< 13
Cladosporium				ND	< 13
Curvularia				ND	< 13
Nigrospora				ND	< 13
Penicillium/Aspergillus types†				ND	< 13
Stachybotrys				ND	< 13
Torula				ND	< 13
Seldom found growing indoors**					
Ascospores††				ND	< 13
Basidiospores††				ND	< 13
Rusts				ND	< 13
Smuts, Periconia, Myxomycetes††				2	27
Total					27

100	MoldSCORE; 100 200 300 Score									
100	200	300	Score							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			100							
			105							
Fina	Final MoldSCORE									

Location: 20804001-TM22JL

Fungi Identified	Indoor sample spores/m3								3	Raw	Spores/	
	<100			١K			10K		>10	0K	count	m3
Generally able to grow indoors*												
Alternaria											ND	< 13
Bipolaris/Drechslera group									Ш		ND	< 13
Chaetomium									Ш		ND	< 13
Cladosporium											1	53
Curvularia									Ш		ND	< 13
Nigrospora									Ш		ND	< 13
Penicillium/Aspergillus types†											ND	< 13
Stachybotrys											ND	< 13
Torula											ND	< 13
Seldom found growing indoors**												
Ascospores††											ND	< 13
Basidiospores††											ND	< 13
Rusts											ND	< 13
Smuts, Periconia, Myxomycetes††											ND	< 13
Total												53

MoldSCORE;									
			100						
			100						
			100						
			101						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
			100						
Fina	101								

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM23JL

Fungi Identified	Iı	ndo	00	rs	sam	ple	e s	por	es/ı	m3	;	Raw	Spores/
	<10	00			1K			10K	2	>100)K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group												ND	< 13
Chaetomium												1	13
Cladosporium												ND	< 13
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												1	53
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores††												ND	< 13
Basidiospores††												ND	< 13
Rusts												ND	< 13
Smuts, Periconia, Myxomycetes††												ND	< 13
Total													66

MoldSCORE‡									
100 200 300	Score								
	100								
	100								
	121								
	100								
	100								
	100								
	106								
	100								
	100								
	100								
	100								
	100								
	100								
Final MoldSCORE	121								

Location: 20804001-TM24JL

Fungi Identified	Indoor sample spores/m3								3	Raw	Spores/
	<100		1K			10 F	(>10	0K	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										1	53
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores††										ND	< 13
Basidiospores††										1	53
Oidium										1	13
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes††										ND	< 13
Total											119

MoldSCORE;									
	Score								
	100								
	100								
	100								
	100								
	100								
	100								
	100								
	100								
	100								
	100								
	105								
	105								
	100								
	100								
Final MoldSCORE	105								

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM25JL

Fungi Identified	Indo	or	sam	ple :	spore	es/n	13	Raw	Spores/
	<100		1K		10K	>	100k	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								1	53
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								2	107
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									160

MoldSCORE: 200 300	Score
	100
	100
	100
	100
	100
	100
	111
	100
	100
	100
	100
	100
	100
Final MoldSCORE	111

Location: 20804001-TM26JL

Fungi Identified	Ind	Indoor sample spores/m3							Spores/
	<100	1	K		10K	>	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								ND	< 13
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								1	53
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									106

MoldSCORE;							
100	Score						
100	100 200 300						
			100				
			100				
			100				
			100				
			100				
			100				
			105				
			100				
			100				
			100				
			105				
			100				
			100				
Fina	al MoldS(CORE	105				

Client: Hygiene Technologies International, Inc.: Northern California

C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM27JI

Fungi Identified	Iı	ıdo	01	rs	samp	ole	S	por	es/i	m3	3	Raw	Spores/
	<10	0			1K			10K		>10	0K	count	m3
Generally able to grow indoors*													
Alternaria												ND	< 13
Bipolaris/Drechslera group												ND	< 13
Chaetomium												ND	< 13
Cladosporium												1	53
Curvularia												ND	< 13
Nigrospora												ND	< 13
Penicillium/Aspergillus types†												36	880
Stachybotrys												ND	< 13
Torula												ND	< 13
Seldom found growing indoors**													
Ascospores††												ND	< 13
Basidiospores††												1	53
Rusts												1	13
Smuts, Periconia, Myxomycetes††												5	67
Total					•								1,066

M HGCODE:								
100 MoldSCORE : 200 300								
	100							
	100							
	100							
	100							
	100							
	100							
	197							
	100							
	100							
	100							
	100							
	104							
	101							
Final MoldSCORE	197							

Location: 20804001-TM28JL

Fungi Identified	Ind	loor s	ampl	Raw	Spores/		
	<100	1	K	10K	>1001	count	m3
Generally able to grow indoors*							
Alternaria						ND	< 13
Bipolaris/Drechslera group						ND	< 13
Chaetomium						ND	< 13
Cladosporium						1	53
Curvularia						ND	< 13
Nigrospora						ND	< 13
Penicillium/Aspergillus types†						10	293
Stachybotrys						ND	< 13
Torula						ND	< 13
Seldom found growing indoors**							
Ascospores††						ND	< 13
Basidiospores††						ND	< 13
Rusts						ND	< 13
Smuts, Periconia, Myxomycetes††						ND	< 13
Total							346

MoldSCOR 200 3	E‡ 00 Score							
	100							
	100							
	100							
	100							
	100							
	100							
	135							
	100							
	100							
	100							
	100							
	100							
	100							
Final MoldSCORI	E 135							

Client: Hygiene Technologies International, Inc.: Northern California

Northern California C/O: Mr. Wes Frey Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM29JL

Fungi Identified	Indo	or s	ample	e s	pore	s/n	13	Raw	Spores/
_	<100	1	K		10K	>1	00K	count	m3
Generally able to grow indoors*									
Alternaria				Ш				ND	< 13
Bipolaris/Drechslera group				Ш				ND	< 13
Chaetomium				Ш				ND	< 13
Cladosporium								2	107
Curvularia				Ш				ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									160

MoldSCORE 100 200 30	E‡ 0 Score
	100
	100
	100
	101
	100
	100
	103
	100
	100
	100
	100
	100
	100
Final MoldSCORE	103

Location: 20804001-TM30JL

Fungi Identified	Indo	or	samj	ple s	spore	es/r	n3	Raw	Spores/
	<100		1K		10K	>	100	count	m3
Generally able to grow indoors*									
Alternaria		Ш		Ш				ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium		Ш						ND	< 13
Cladosporium								3	160
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								ND	< 13
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								ND	< 13
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								2	27
Total									187

100	Score		
			100
			100
			100
			103
			100
			100
			100
			100
			100
			100
			100
			100
			103
Fina	al MoldSC	ORE	103

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey

Re: 20804001

Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

MoldSCORETM: Spore Trap Report

Location: 20804001-TM31JL

Fungi Identified	Inc	Indoor sample spores/m3								Raw	Spores/
	<100		1	K			10K	3	>100	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										ND	< 13
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores††										ND	< 13
Basidiospores††										ND	< 13
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes††										ND	< 13
Total				•				•			N/A

MoldSCORE; 100 200 300 Score						
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
	100					
Final MoldSCORE	100					

Location: 20804001-TM32JL

Fungi Identified	Ind	oor s	ampl	e spor	es/m3	Raw	Spores/
	<100	1	K	10K	>1001	count	m3
Generally able to grow indoors*							
Alternaria						ND	< 13
Bipolaris/Drechslera group						ND	< 13
Chaetomium						ND	< 13
Cladosporium						ND	< 13
Curvularia						ND	< 13
Nigrospora						ND	< 13
Penicillium/Aspergillus types†						1	53
Stachybotrys						ND	< 13
Torula						ND	< 13
Seldom found growing indoors**							
Ascospores††						ND	< 13
Basidiospores††						1	53
Rusts						ND	< 13
Smuts, Periconia, Myxomycetes††						1	13
Total							119

100	MoldSCORE; Score						
			100				
			100				
			100				
			100				
			100				
			100				
			104				
			100				
			100				
			100				
			105				
			100				
			101				
Final	MoldSCO	ORE	105				

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Date of Receipt: 04-15-2008 Date of Report: 04-17-2008

Date of Sampling: 04-14-2008

Re: 20804001

MoldSCORETM: Spore Trap Report

Location: 20804001-TM33JL

Fungi Identified	Indo	or	sam	ple	n3	Raw	Spores/		
	<100		1K		10K	>	100K	count	m3
Generally able to grow indoors*									
Alternaria								ND	< 13
Bipolaris/Drechslera group								ND	< 13
Chaetomium								ND	< 13
Cladosporium								1	53
Curvularia								ND	< 13
Nigrospora								ND	< 13
Penicillium/Aspergillus types†								1	53
Stachybotrys								ND	< 13
Torula								ND	< 13
Seldom found growing indoors**									
Ascospores††								ND	< 13
Basidiospores††								3	160
Rusts								ND	< 13
Smuts, Periconia, Myxomycetes††								ND	< 13
Total									266

MoldSCORE‡										
100	200 300	Score								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		100								
		115								
		100								
		100								
Fina	al MoldSCORE	115								
	I mai Molascotte 113									

^{*}The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.

^{**}These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

[†]The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

^{††}Most of these spore types are not seen with culturable methods (Anderson sampling), although some may appear as non-sporulating fungi. Most of the basidiospores are "mushroom" spores.

[‡]Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.



Report for:

Mr. Wes Frey Hygiene Technologies International, Inc.: Northern California 3127 Bowen Island Street West Sacramento, CA 95691

Regarding: Project: 20804001

EMĹ ID: 410940

Approved by:

Lab Manager Magzoub Ismail Dates of Analysis: Culturable air fungi (Incl. Asp spp.): 04-22-2008 Spore trap analysis: 04-17-2008

Project SOPs: Culturable air fungi (Incl. Asp spp.) (I100002), Spore trap analysis (I100000)

This coversheet is included with your report in order to comply with AIHA and ISO accreditation requirements.

For clarity, we report the number of significant digits as calculated; but, due to the nature of this type of biological data, the number of significant digits that is used for interpretation should generally be one or two. All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank corrections of results is not a standard practice. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

Document Number: 200091 - Revision Number: 5

5473 Kearny Villa Road, Suite 130, San Diego, CA 92123 (858) 569-5800 Fax (858) 569-5806 www.emlab.com

Client: Hygiene Technologies International, Inc.:

Northern California C/O: Mr. Wes Frey Re: 20804001 Date of Sampling: 04-14-2008 Date of Receipt: 04-15-2008 Date of Report: 04-22-2008

CULTURABLE AIR FUNGI REPORT

Location:		04001-)1outJL	2080400	1-VM02JL	20804001-VM03JL		2080400	1-VM04JL
Comments (see below)	N	Vone	N	lone	N	lone	None	
Lab ID-Version‡:	180	5499-1	180:	5500-1	180	5501-1	180:	5502-1
	raw ct.	cfu*/m3	raw ct.	cfu*/m3	raw ct.	cfu*/m3	raw ct.	cfu*/m3
Acremonium								
Alternaria	1	18						
Aphanocladium								
Arthrospore-former			1	18				
Aspergillus flavus								
Aspergillus fumigatus								
Aspergillus nidulans								
Aspergillus niger								
Aspergillus ochraceus								
Aspergillus versicolor								
Aureobasidium								
Basidiomycetes								
Bipolaris/Drechslera group								
Botrytis								
Chaetomium	1	18						
Cladosporium	60	1,150			1	18	1	18
Curvularia								
Epicoccum								
Fusarium								
Non-sporulating fungi	2	35						
Paecilomyces								
Penicillium								
Phoma								
Rhizopus								
Stachybotrys chartarum								
Ulocladium								
Yeasts	1	18						
Positive Hole	400	_	400		400		400	
Sample volume (liters)	56.6		56.6		56.6		56.6	
TOTAL CFU*/M3		1,239		18		18		18

^{*} cfu = colony forming units

Positive hole correction chart used for all calculations

Comments:

Note: Interpretation is left to the company and/or persons who conducted the field work. Variation is an inherent part of biological sampling. The presence or absence of a few genera in small numbers should not be considered abnormal.

NORMAL SPORE LEVELS: Indoor spore levels usually average 30 to 80% of the outdoor spore level at the time of sampling, with the same general distribution of spore types. Filtered air, air-conditioned air, or air remote from outside sources may average 5 to 15% of the outside air at the time of sampling. (These percentages are guidelines, only. A major factor is the accessibility of outdoor air. A residence with open doors and windows and heavy foot traffic may average 95% of the outdoor level while high rise office buildings with little air exchange may average 2%. Dusty interiors may exceed 100% of the outdoors to some degree, but will still mirror the outdoor distribution of spore types.)

PROBLEM INTERIORS: A substantial increase of one or two spore types which are inconsistent with and non-reflective of the outside distribution of spore types is usually indicative of an indoor reservoir of mold growth.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

5473 Kearny Villa Road, Suite 130, San Diego, CA 92123 (858) 569-5800 Fax (858) 569-5806 www.emlab.com

Client: Hygiene Technologies International, Inc.:

Northern California

C/O: Mr. Wes Frey

Date of Sampling: 04-14-2008

Date of Receipt: 04-15-2008

Date of Report: 04-22-2008

Re: 20804001

CULTURABLE AIR FUNGI REPORT

Location:	2080400	1-VM05JL	2080400	1-VM06JL	2080400	1-VM07JL	2080400	1-VM08JL
Comments (see below)	N	lone	N	lone	N	lone	N	lone
Lab ID-Version‡:	180:	5503-1	1803	5504-1	180	5505-1	180	5506-1
	raw ct.	cfu*/m3						
Acremonium								
Alternaria					1	18		
Aphanocladium								
Arthrospore-former								
Aspergillus flavus								
Aspergillus fumigatus								
Aspergillus nidulans								
Aspergillus niger								
Aspergillus ochraceus								
Aspergillus versicolor								
Aureobasidium								
Basidiomycetes								
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium					2	35	1	18
Curvularia								
Epicoccum								
Fusarium								
Non-sporulating fungi								
Paecilomyces								
Penicillium	1	18	2	35				
Phoma								
Rhizopus								
Stachybotrys chartarum								
Ulocladium								
Yeasts			1	18				
Positive Hole	400		400		400		400	
Sample volume (liters)	56.6		56.6		56.6		56.6	
TOTAL CFU*/M3	1 1	18	1 . 1	53		53		18

^{*} cfu = colony forming units

Positive hole correction chart used for all calculations

Comments:

Note: Interpretation is left to the company and/or persons who conducted the field work. Variation is an inherent part of biological sampling. The presence or absence of a few genera in small numbers should not be considered abnormal.

NORMAL SPORE LEVELS: Indoor spore levels usually average 30 to 80% of the outdoor spore level at the time of sampling, with the same general distribution of spore types. Filtered air, air-conditioned air, or air remote from outside sources may average 5 to 15% of the outside air at the time of sampling. (These percentages are guidelines, only. A major factor is the accessibility of outdoor air. A residence with open doors and windows and heavy foot traffic may average 95% of the outdoor level while high rise office buildings with little air exchange may average 2%. Dusty interiors may exceed 100% of the outdoors to some degree, but will still mirror the outdoor distribution of spore types.)

PROBLEM INTERIORS: A substantial increase of one or two spore types which are inconsistent with and non-reflective of the outside distribution of spore types is usually indicative of an indoor reservoir of mold growth.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

5473 Kearny Villa Road, Suite 130, San Diego, CA 92123 (858) 569-5800 Fax (858) 569-5806 www.emlab.com

Client: Hygiene Technologies International, Inc.:

Northern California

C/O: Mr. Wes Frey

Date of Sampling: 04-14-2008

Date of Receipt: 04-15-2008

Date of Report: 04-22-2008

Re: 20804001

CULTURABLE AIR FUNGI REPORT

Location:	2080400	1-VM09JL	20804001-VM10outJL			
Comments (see below)	N	Vone	None			
Lab ID-Version‡:	180	5507-1	1805508-1			
	raw ct.	cfu*/m3	raw ct.	cfu*/m3		
Acremonium						
Alternaria			1	18		
Aphanocladium			1	18		
Arthrospore-former						
Aspergillus flavus						
Aspergillus fumigatus	1	18				
Aspergillus nidulans			1	18		
Aspergillus niger			1	18		
Aspergillus ochraceus						
Aspergillus versicolor						
Aureobasidium			2	35		
Basidiomycetes						
Bipolaris/Drechslera group						
Botrytis						
Chaetomium						
Cladosporium	1	18	71	1,380		
Curvularia				,		
Epicoccum						
Fusarium						
Non-sporulating fungi	1	18				
Paecilomyces Paecilomyces						
Penicillium	1	18	3	53		
Phoma						
Rhizopus						
Stachybotrys chartarum						
Ulocladium						
Yeasts	1	18	3	53		
Positive Hole	400		400			
Sample volume (liters)	56.6		56.6			
TOTAL CFU*/M3		90		1,593		

^{*} cfu = colony forming units

Positive hole correction chart used for all calculations

Comments:

Note: Interpretation is left to the company and/or persons who conducted the field work. Variation is an inherent part of biological sampling. The presence or absence of a few genera in small numbers should not be considered abnormal.

NORMAL SPORE LEVELS: Indoor spore levels usually average 30 to 80% of the outdoor spore level at the time of sampling, with the same general distribution of spore types. Filtered air, air-conditioned air, or air remote from outside sources may average 5 to 15% of the outside air at the time of sampling. (These percentages are guidelines, only. A major factor is the accessibility of outdoor air. A residence with open doors and windows and heavy foot traffic may average 95% of the outdoor level while high rise office buildings with little air exchange may average 2%. Dusty interiors may exceed 100% of the outdoors to some degree, but will still mirror the outdoor distribution of spore types.)

PROBLEM INTERIORS: A substantial increase of one or two spore types which are inconsistent with and non-reflective of the outside distribution of spore types is usually indicative of an indoor reservoir of mold growth.

The Limit of Detection is the product of a raw count of I and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.



Hygiene Technologies International, Inc.

3625 Oct Ama Boulevard, Suite 180 Torrance, Cal fomia 90503-1643 (310) 370-8370

(\$10) 370-2474 FAX www.hygienetech.com

					
Project Number/Purcha	se Order: 20	804001	Date \$	Submitted: <u>4</u> /	14/08
Project Contact:			Turnaround Requi		
Lab Destination:		u	Lab Contact:		
SAMPLE ID	VOLUME	MEDIA	ANALYSI	S REQUESTED	
20804001 - VM 0104175L		MEA	vialde fungi	··	
- VM02-31		1			
-VM037L					
- AWU4ar		ļ		· <u>- :</u>	
-VM053L	<u> </u>				·
-VW962F		·			
-MOTTL		ļ . ļ			
- VMO82F			ļ		
-VM095L			ļi		
-VM100UT3	1	W	<u> </u>	···	<u>. </u>
-TMI TOUT 3L	751	allergenco >	total fungi		
-TM182L		 	1	•	
TMIAZE			· · ·		
-TMZ03L	 				·
-TM2131		$+$ \downarrow	-		
Special Instructions:		1	<u>'</u> ¥		
Preciae misis actività.					• .
		:			
1. Sampled by:	holombe Ali	i.las 1600	Received by:	MORE 4	ISING IDAY
2. Relinquished by			Received by:		μ <i>ν(να∠\</i> ν ε% .
3. Relinquished by					
		Please include sig	mature, date, and time		
Lab Use Only:				-	<u> </u>
					אָןוּטִיאָרֶ
1					L



Hygiene Technologies International, Inc.

3625 Del Amo Roulevard, Suite 180 Torrance, Galifornia 90503-1643 (310) 370-8370 (310) 370-2474 FAX

www.hygienetech.com

					
Project Number/Purcha	ise Order:	L0804001		Date Submi	tted: 4/14/08
Project Contact:	Wes Frey	}	Turna	round Required: _	Standard
Lab Destination:	<u>eml</u>		Lab C	ontact;	
SAMPLE ID	VOLUME	MEDIA	<u>.</u>	ANALYSIS REC	QUESTED
20804001-TM035L	751	allergenco.D	total	Lungi ID	
-TM 24-51	1 .				
-TM253L	\ <u>\</u>	<u> </u>		·	
-7M203L	<u> </u>				
-1M273L	<u> </u>	· · · · · · · · · · · · · · · · · · ·			
-TM 2.932	 	<u> </u>		<u> </u>	
-7M293L			<u> </u>		<u>·</u>
-144 302F					
		 	·	- 	· -
-IW752F		 	<u> </u>		
-TM3331	1 1/			\downarrow	
TM 34 out	<u> </u>	1			
					-
		·			
	-				
Special Instructions	·				.
			· 		
1. Sampled by:	John Le 41	14/08 1600	Received 1	by: SAXXAIT	E86 4/15/08 10A
2. Relinquished by	•	' '		by:	I
3. Relinquished by	/:	·	Received 1	by:	
		Please include sig	nature, date, and	d time	
Lab Use Only:					
					(
<u> </u>					

Client: Hygiene Technologies International, Inc.

C/O: Mr. John Le Date of Receipt: 04-11-2008 Re: 20804001 Date of Report: 04-14-2008

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		04001- 9CCJL		04001- 0CCJL	20804001- TM11CCJL		20804001- TM12CCJL	
Comments (see below)	N	lone	N	lone	N	Vone	None	
Lab ID-Version‡:	180	1683-1	180	1684-1	180	1685-1	180	1686-1
	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3	raw ct.	spores/m3
Alternaria								
Arthrinium								
Ascospores*								
Aureobasidium								
Basidiospores*							1	53
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium	2	107	1	53	2	107	1	53
Curvularia								
Epicoccum					1	13		
Fusarium								
Myrothecium								
Nigrospora								
Other colorless								
Penicillium/Aspergillus types†			2	107				
Pithomyces								
Rusts*								
Smuts*, Periconia, Myxomycetes*								
Stachybotrys								
Stemphylium								
Torula								
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	1+		2+		2+		2+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		107		160		120		106

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens. † The spores of *Aspergillus* and *Penicillium* (and others such as *Acremonium*, *Paecilomyces*) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

Client: Hygiene Technologies International, Inc.

C/O: Mr. John Le Date of Receipt: 04-11-2008 Re: 20804001 Date of Report: 04-14-2008

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:		04001- 3CCJL		04001- 4CCJL		04001- 5CCJL		04001- 6CCJL
Comments (see below)	N	lone	N	lone	N	Vone	N	lone
Lab ID-Version‡:	180	1687-1	180	1688-1	180	1689-1	180	1690-1
	raw ct.	spores/m3						
Alternaria				_		_		_
Arthrinium								
Ascospores*								
Aureobasidium								
Basidiospores*								
Bipolaris/Drechslera group								
Botrytis								
Chaetomium								
Cladosporium	1	53	4	213			3	160
Curvularia								
Epicoccum								
Fusarium								
Myrothecium								
Nigrospora								
Other colorless								
Penicillium/Aspergillus types†			1	53	2	107		
Pithomyces								
Rusts*								
Smuts*, Periconia, Myxomycetes*	5	67	2	27				
Stachybotrys								
Stemphylium								
Torula			1	13				
Ulocladium								
Zygomycetes								
Background debris (1-4+)††	3+		3+		3+		3+	
Sample volume (liters)	75		75		75		75	
TOTAL SPORE/m3		120		306		107		160

Comments:

^{*} Most of these spore types are not seen with culturable methods (Andersen sampling), although some may appear as non-sporulating fungi.

Most of the basidiospores are "mushroom" spores while the rusts and smuts are plant pathogens. † The spores of *Aspergillus* and *Penicillium* (and others such as *Acremonium*, *Paecilomyces*) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher then reported. It is important to account for samples volumes when evaluating dust levels.

The Limit of Detection is the product of a raw count of 1 and 100 divided by the percent read. The analytical sensitivity (counts/m3) is the product of the Limit of Detection and 1000 divided by the sample volume.

[‡] A "Version" greater than 1 indicates amended data.

EMLab P&K

3800 Watt Ave., Suite 145, Sacramento, CA 95821 (650) 829-5800 Fax (650) 829-5852 www.emlab.com

Client: Hygiene Technologies International, Inc. C/O: Mr. John Le

Re: 20804001

Date of Sampling: 04-10-2008 Date of Receipt: 04-11-2008 Date of Report: 04-14-2008

DIRECT MICROSCOPIC EXAMINATION REPORT

(Wet Mount)

(
Background Debris and/or Description	Miscellaneous Spores Present*	MOLD GROWTH: Molds seen with underlying mycelial and/or sporulating structures†	Other Comments††	General Impression
Lab ID-Version‡: 1	801667-1: Swab sar	mple 20804001-S01JL		
Moderate	Few	None	None	Normal trapping
Lab ID-Version: 18	801668-1: Swab sam	nple 20804001-S02JL		
Heavy	Few	None	Many <i>Ulocladium</i> spores detected.	Mold growth in vicinity?
Lab ID-Version: 18	01669-1: Swab sam	ple 20804001-S03JL		
Heavy	Few	None	Many <i>Ulocladium</i> and <i>Cladosporium</i> spores detected.	Mold growth in vicinity?
Lab ID-Version: 18	301670-1: Swab sam	ple 20804001-S04JL		
Heavy	Few	None	Many <i>Ulocladium</i> and colorless spores typical of <i>Penicillium / Aspergillus</i> detected.	Mold growth in vicinity?
Lab ID-Version: 18	301671-1: Swab sam	nple 20804001-S05JL		
Heavy	Few	None	Many colorless spores typical of Penicillium / Aspergillus detected.	Mold growth in vicinity?
Lab ID-Version: 18	301672-1: Swab sam	pple 20804001-S06JL		
Heavy	Few	None	Many colorless spores typical of Penicillium / Aspergillus detected.	Mold growth in vicinity?
Lab ID-Version: 18	01673-1: Swab sam	pple 20804001-S07JL		
Moderate	Few	None	None	Normal trapping
Lab ID-Version: 18	301674-1: Swab sam	nple 20804001-S08JL	1	
Heavy	Few	None	Many <i>Alternaria</i> spores detected.	Mold growth in vicinity?
Lab ID-Version: 18	801675-1: Swab sam	nple 20804001-S09JL		
Moderate	Few	None	None	Normal trapping
			EMLab	ID: 410145, Page 1 of 2

Background Debris and/or	Miscellaneous	MOLD GROWTH: Molds seen	Other Comments††	General				
Description	Spores Present*	with underlying mycelial and/or sporulating structures†	Comments	Impression				
	1801676-1: Swab sa	mple 20804001-S10JL						
Heavy	Few	None	None	Normal trapping				
Lab ID-Version: 18	Lab ID-Version: 1801677-1: Swab sample 20804001-S11JL							
Moderate	Few	None	None	Normal trapping				
1vioderate	10,,		Tione	Tronnar trapping				
Lab ID-Version: 18		pple 20804001-S12JL	<u> </u>	<u> </u>				
Moderate	Few	None	None	Normal trapping				
Lab ID-Version: 18	301679-1: Swab sam	pple 20804001-S13JL						
Heavy	Few	None	None	Normal trapping				
Lab ID-Version: 18	 801680-1: Swab sam	pple 20804001-S14JL						
Heavy	Very few	None	None	Normal trapping				
Lab ID-Version: 18	301681-1: Swab sam	ple 20804001-S15JL						
Heavy	Few	None	None	Normal trapping				
X 1 XD XX : 10	201602.1.6.1	1. 2000 4001 (316)						
	801682-1: Swab sam	pple 20804001-S16JL	ı					
Moderate	Few	None	Many <i>Alternaria</i> spores detected.	Mold growth in vicinity?				
† A "Vargion" greater th	4 . 1	1 .	I .					

[‡] A "Version" greater than 1 indicates amended data.



Hygiene Technologies International, Inc.

3625 Del Amo Soulevard, Suite 180 Torrence, Californie 90503-1643

(310) 370-8370 (310) 370-2474 FAX www.hyglenetech.com

Project Number/Purcha	se Order:	280400	Date Submitted:	4/10/08
Project Contact:	Wes Fra	}	Turnaround Required:	undard
Lab Destination:	EM lab	···-	Lab Contact;	
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUEST	
20804001-809JL	N/A-	Swaln	surface fungi ID qu	relitative
-8105L	. 1	· · · · · ·	0 0	
		<u> </u>		<u> </u>
- 81 2 JL	· ·	<u> </u>		
-81372		· · · · · · · · · · · · · · · · · · ·		
<u>-8143L</u>	<u> </u>			
-21627				<u> </u>
-6163L	V	1	1 1 1 2	
-TMORCCEL	1	lallergenco 1/	total fungi ID	
-TMIOCCOL	· · · · · · · · · · · · · · · · · · ·			·
-TMIRCOIL	 -			
-TMIRCOSI		 - 		·
-TM14CCJL	<u> </u>			
-TAUSCC3L				
-TMIGCCOL	1 1/			
Special Instructions:				
1. Sampled by:	ohnle 4	10/08 1200	Received by: Ossanlen Cooley	0\$/11/08 400pm
2. Relinquished by:			Received by:	
3. Relinquished by:	* .	·	Received by:	
· · · · · · · · · · · · · · · · · · ·		Please include sig	nature, date, and time	
Lab Use Only:			•	401-R
				<u>-</u>
			·	91



Hygiene Technologies International, Inc.

3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643 (310) 370-8370

> 310) 370-2474 FAX www.hyglenetech.com

· 			
Project Number/Purcha	ise Order:	0804001	Date Submitted: 4/11/08
Project Contact:	Wes Fr	reur	Turnaround Required:andard
Lab Destination:	EM lalos	<u> </u>	Lab Contact:
SAMPLE ID	VOLUMB	MEDIA	ANALYSIS REQUESTED
20804001- SOIJL	N/A	Swah	surface Jungi ID qualitative
- S0231	1 1		· · · · · · · · · · · · · · · · · · ·
- S035L		·	<u> </u>
-50431	<u> </u>	<u> </u>	
- SOSTL			
-50671			<u> </u>
-8075L	1 1/	17:	
- 50831	· · · · · · · · · · · · · · · · · · ·	· V	· · · · · · · · · · · · · · · · · · ·
· · · · · · · · · · · · · · · · · · ·		 	
· ·	· .		
		·	
			-
·	<u> </u>		
-	.1	······································	
Special Instructions:			
·			7 1 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1. Sampled by:	• • •	08 1500_	Received by: Cassandra Cooling 11/11/08 400pm
2. Relinquished by:		•	Received by:
3. Relinquished by:		Please include sig	Received by:
Lab Use Only:			
			7. F.
			2



Page 1 of 6

2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

FINAL REPORT: Total Fungal Spore Trap Count

PROJECT NUMBER: 20804001

LABORATORY ID NUMBER: 0804013

Hygiene Technologies International, Inc.

Received Date:

April 07, 2008

Attention: Wes Frey

Report Date:

4330 Auburn Blvd. Suite 1850

April 11, 2008

Sacramento, CA 95841

Customer Sample Number: -TM01CCJL Background: Moderate particulates	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 11-Apr-08	Detection Limit:	52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Ascospores			Р	
Basidiospores		1	52	
Cladosporium		4	210	
Fusarium			P	
Oidium		1	52	
Penicillium/Aspergillus types		3	160	
Pollen			P	
Smuts/ Myxomycetes		1	52	
Stachybotrys			P	
Unidentified mitosporic fungi			P	
TOTAL		10	530	

P - Spores Present

« (less than) = measurement below the reporting limit

Rounding: Note that all reported counts have been rounded to two significant figures based on the sampling and analytical methods used. BioHygiene Labs rounds such that if the last significant digit is an even number, then the result is rounded down to that digit; if the last significant digit is an odd number, then it is rounded up to the nearest even number. Thus the TOTAL may not equal the sum of the individual counts per column. TOTAL rows do not include pallen.

Background is graded as Very Light (0 - 10%), Light (>10 - 30%), Moderate (>30 - 70%), Heavy (>70 - 90%), and Very Heavy (>90%) Particulates as a percentage of the trace area.

DATE: 04/11/08

Name

Title:

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.



Page 2 of 6

2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

FINAL REPORT: Total Fungal Spore Trap Count

PROJECT NUMBER: 20804001

LABORATORY ID NUMBER: 0804013

Hygiene Technologies International, Inc.

Received Date:

April 07, 2008

Report Date:

April 11, 2008

Attention: Wes Frey 4330 Auburn Blvd. Suite 1850 Sacramento, CA 95841

Customer Sample Number: -TM02CCJL Background: Heavy particulates	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 11-Apr-08	Detection Limit:	52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Alternaria		2	100	
Ascaspores		3	160	
Basidiospores		13	680	
Botrytis			P	
Cladosporium		57	3000	
Penicillium/ Aspergillus types		5	260	
Pollen		3	160	
Rusts			P	
Smuts/Myxomycetes		5	260	
Torula		- 4	52	
Ulocladium			P	
Unidentified mitosporic fungi			P	
TOTAL		86	4500	

P = Spores Present

< (less than) = measurement below the reporting limit

Rounding: Note that all reported counts have been rounded to two significant figures based on the sampling and analytical methods used. BioHygiene Labs rounds such that if the last significant digit is an even number, then the result is rounded down to that digit; if the last significant digit is an odd number, then it is rounded up to the nearest even number. Thus the TOTAL may not equal the sum of the individual counts per column. TOTAL rows do not include pollen.

Background is graded as Very Light (0 - 10%), Light (>10 - 30%), Moderate (>30 - 70%), Heavy (>70 - 90%), and Very Heavy (>90%) Particulates as a percentage of the trace area.

APPROVED:	1600		DATE:	04/11/08	
Nameluc	as Wallin	Title:	Ind Amelyst		

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.



2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

Page 3 of 6

FINAL REPORT: Total Fungal Spore Trap Count

PROJECT NUMBER: 20804001

LABORATORY ID NUMBER: 0804013

Received Date:

Report Date:

Hygiene Technologies International, Inc.

April 07, 2008

Attention: Wes Frey

4330 Auburn Blvd. Suite 1850

April 11, 2008

Sacramento, CA 95841

Customer Sample Number: -TM03CCJL Met	-d- 11101 1	400000000000000000000000000000000000000	
	od: M101.1 Date Of Analysis: 11-Apr-08 le Intact: Yes	Detection Limit:	52 Spores/M ³
Genus (species)	Raw Count	Total Spores / M ³	Comment
Alternaria	2	100	
Ascospores	1	52	
Basidiospores	9	470	
Botrytis		Р	
Cladosporium	39	2000	
Nigrospora	1014	Р	
Oidium		. P	
Penicillium/Aspergillus types	3	160	
Pollen	3	160	
Rusts		P	
Smuts/Myxomycetes	4	210	
Stemphylium		P	
Ulocladium		P	
Unidentified mitosporic fungi	4	210	
TOTAL	62	3200	

P = Spores Present

< (less than) - measurement below the reporting limit

Rounding: Note that all reported counts have been rounded to two significant figures based on the sampling and analytical methods used. BioHyglene Labs rounds such that if the last significant digit is an even number, then the result is rounded down to that digit; if the last significant digit is an odd number, then it is rounded up to the nearest even number. Thus the TOTAL may not equal the sum of the individual counts per column. TOTAL rows do not include pollen.

Background is graded as Very Light (0 - 10%), Light (>10 - 30%), Moderate (>30 - 70%), Heavy (>70 - 90%), and Very Heavy (>90%) Particulates as a percentage of the trace area.

APPROV	ED:	7	DATE: 04/11/08	
Name _	Lucas Wallen	Title:	Lab Analyst	

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.



Page 4 of 6

2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

FINAL REPORT: Total Fungal Spore Trap Count

PROJECT NUMBER: 20804001		L	ABORATORY ID NUMB	ER: 0804013
Hygiene Technologies Internationa	l, Inc.		Received Date:	April 07, 2008
Attention: Wes Frey			Report Date:	April 11, 2008
4330 Auburn Blvd. Suite 1850			report bate.	April 11, 2006
Sacramento, CA 95841				
Customer Sample Number: -TM04CCJL Background: Moderate particulates	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 11-Apr-08	Detection Limit	: 52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Alternaria			Р	
Ascospores		1	52	
Basidiospores		2	100	
Cladosporium		3	160	
Penicillium/Aspergillus types		5	260	
Smuts/Myxomycetes			Р	
Unidentified mitosporic fungi		3	160	
Unidentified zygomycetes			Р	
TOTAL		14	730	
Customer Sample Number: -TM05CCJL Background: Moderate particulates	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 10-Apr-08	Detection Limit	: 52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Ascospores		1	52	
Basidiospores		5	260	
Cladosporium		3	160	
Epicoccum			P	
Penicillium/Aspergillus types		2	100	
Pollen			P	
Smuts/Myxomycetes		2	100	
Torula		1	52	
Unidentified zygomycetes		1	52	
TOTAL		15	780	
P = Spores Present < (less than) = measures Rounding: Note that all reported counts have been rounded digit is an even number, then the result is rounded down to equal the sum of the individual counts per column. TOTAL Background is graded as Very Light (0 · 10%), Light (>10 · 30%)	that digit; if the last significant or rows do not include pollen.	on the sampling and analytical methods used ligit is an odd number, then it is rounded up	to the nearest even number. Thus th	e TOTAL may not
APPROVED:	<u>'</u>	DATE: Du///	(vB	

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.



Page 5 of 6

2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

April 11, 2008

Report Date:

FINAL REPORT: Total Fungal Spore Trap Count

PROJECT NUMBER: 20804001

LABORATORY ID NUMBER: 0804013 Hygiene Technologies International, Inc. Received Date: April 07, 2008

Attention: Wes Frey

4330 Auburn Blvd. Suite 1850

Sacramento, CA 95841 Customer Sample Number: -TM06CCJL Method: M101.1

Background: Moderate particulates	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 10-Apr-08	Detection Limit:	52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Basidiospores		4	. Ctal Spores / M	Comment
Cladosporium		<u> </u>	52	
Penicillium/Aspergillus types		1	52	
Smuts/Myxomycetes		2	100	
Unidentified mitosporic fungi			Р	
TOTAL			Р	•
TOTAL		4	200	

Customer Sample Number: -TM07CCJL Background: Moderate particulates Genus (species)	Method: M101.1 Sample Intact: Yes	Date Of Analysis: 11-Apr-08	Detection Limit:	52 Spores/M ³	
Alternaria		Raw Count	Total Spores / M ³	Comment	
			P		
Ascospores			<u> </u>		
Basidiospores			P		
Cladosporium		5	260		
Oidium		2	100		
Penicillium/Aspergillus types			P		
Pollen		7	360		
Rusts			Р		
Smuts/Myxomycetes			P		
		2	100		
Unidentified mitosporic fungi					
TOTAL			P		
		16	820		

P = Spores Present

Rounding: Note that all reported counts have been rounded to two significant figures based on the sampling and analytical methods used. BioHygiene Labs rounds such that if the last significant digit is an even number, then the result is rounded down to that digit; if the last significant digit is an odd number, then it is rounded up to the nearest even number. Thus the TOTAL may not

Background is graded as Very Light (0 - 10%), Light (>10 - 30%), Moderate (>30 - 70%), Heavy (>70 - 90%), and Very Heavy (>90%) Particulates as a percentage of the trace area.

APPROVED:	DATE:64/u/06
Name Lun (Wallin Title:	Lat Analyst
Results reported relate only to the sample it an	

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.

< (less than) = measurement below the reporting limit



PROJECT NUMBER: 20804001

Hygiene Technologies International, Inc.

Page 6 of 6

TOTAL

Attention: Wes Frey

4330 Auburn Blvd. Suite 1850 Sacramento, CA 95841

2200 Amapola Ct., Suite 102 Torrance, CA 90501 (310) 618-0400 (310) 618-9400 fax

FINAL REPORT: Total Fungal Spore Trap Count

LABORATORY ID NUMBER: 0804013

200

Received Date:

April 07, 2008

Report Date:

April 11, 2008

Customer Sample Number: -TM08CCJL Background: Light particulates	Method: M101,1 Sample Intact: Yes	Date Of Analysis: 11-Apr-08	Detection Limit:	52 Spores/M ³
Genus (species)		Raw Count	Total Spores / M ³	Comment
Basidlospores		2	100	
Botrytis			P	
Cladosporium		2	100	
Unidentified mitosporic fungi			P	

4

P « Spores Present

< (less than) - measurement below the reporting limit

Rounding: Note that all reported counts have been rounded to two significant figures based on the sampling and analytical methods used. BioHygiene Labs rounds such that if the last significant digit is an even number, then the result is rounded down to that digit; if the last significant digit is an odd number, then it is rounded up to the nearest even number. Thus the TOTAL may not equal the sum of the individual counts per column. TOTAL rows do not include pollen.

Background is graded as Very Light (0 + 10%), Light (>10 - 30%), Moderate (>30 - 70%), Heavy (>70 - 90%), and Very Heavy (>90%) Particulates as a percentage of the trace area.

APPROVE	ED:	4		DATE:	04/11/08	
Name	lucas	Willia	Title:	lab Analyst		

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.



-0804013

Hygiene Technologies International, Inc.

3625 Del Amo Boulevard, Sulte 180 Torrance, California 90503-1643 (310) 370-8370 (310) 370-2474 FAX

Project Number/Purchas	se Order:	20804001	Date Submitted: 4/4/08
Project Contact:			Turnaround Required: Standard
Lab Destination:	Bio Hyg	iene	Lab Contact:
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED
20804001 - TMOICCOL	75L	allergenco D	N 101.1
20804001 - TMODECEL			
20804001 - TM03CCSL		+	
20804001 - TMOACCSL			
20804001 - TMOSCOTE			
20804001 - TMUGCCSL			
20804001 - TMOTECSI			
20804001-TMORCUSE	W	V	V
		MOY O	202
			,
*	- a		+
Special Instructions:			
1. Sampled by:	ohn Le 4	14/08 1500	Received by: Pallemoy 03 9:36
			Received by: 12 Sulleys 04.19.08 9:10-Thorace
3. Relinquished by:	70		
Con The Month Process II Bases		Please include signa	thre, date, and time UNW 0410-29 16:30 - THORES THOSE
Lab Use Only: Contac	ted Tomile ou	10.00 to con form o	CCTL, - Thoracute) p8 (- Mod cute) p11 (-TMBCC)
impleted as our	1.08 800F S	~ 1 P + (-Imu)	(CC)2,-morcusz) p8(-mov(cusz) p11(-mBcc)
- N. I	professor agency agency		
Completed on ou	mus Berk 33	104 p.7 (-TM074)	st, -tmosccze). ww

3

5

Hygiene Technologies International, Inc.

-0804013

10/23

3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643 (310) 370-8370 (310) 370-2474 FAX

www.hygienetech.com

Project Number/Purcha	se Order:	20804001			
Project Contact;	Wes Fr	ey	Turnaround Required: Standard		
Lab Destination:	Bio Hygie	ne	Lab Contact:		
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED		
20804001-TLOIJL	N/A	Bio tape	M 102.1		
20804001 - TLOUGL					
20804001 - TL035L		100			
20804001 - TLOAJL					
10804001 - TLOSJL					
20804001 -TLOGSL			/2		
20804001 - TLO75L					
20804001 -TLOSTL		+			
20804001 - TL09JL					
20804001 -TL105L					
20804001 - TLIIJL					
2080400 - TLIZJL					
20804001 -TL135L		12			
20804001 -TLI45L					
2.0804001 TLISTL					
20804001 - TLIGSL	A	1 4			
Special Instructions:					
			*		
			1-A 110		
1. Sampled by:	1040101		Received by: Paulleys 01 07-08 9:30		
2. Relinquished by:	Halley	704.02.08/0	Preceived by: R. Dalley 04.10.08 71/40-12/6		
	0		Received by:		
3. Relinquished by:					



Hygiene Technologies International, Inc.

2017-3

3625 Del Amo Boulevard, Suite 180 Torrance, California 90503-1643

> (310) 370-8370 (310) 370-2474 FAX

Project Number/Purchas	se Order:	20904001	Date Submitted: 4/4/08
Project Contact:	Wes Frey	4	Turnaround Required: Standard
Lab Destination:	The state of the s	A .	Lab Contact:
SAMPLE ID	VOLUME	MEDIA	ANALYSIS REQUESTED
20804001 - TL175L	NA	Bio tape	1. co M
20804001 - TL18JL		1	
20804001 -TL195L		-	
20804001 - TL 205L	V	V	
		ten	
		1	Page .
		ii ii	
2			
Special Instructions:			
			1000
1. Sampled by:	ohyle 4	14/08 1200	Received by: Pha Close 04 07-08 7.39
2. Relinquished by:	Hung	704.07.08 10:3	Received by: Poulleyer 04.11.08 9:10
3. Relinquished by:		Please include sid	Received by:
Lab Use Only	-		Control of the Contro
Completed in our 11.	08 BOOK 32	02 p21-12173	2) p 10(-11852+0-12052)e6.



2200 Amapola Ct., Suite 102 Torrance, CA 90501

(310) 618-0400 (310) 618-9400 fax

Page 1 of 3

FINAL REPORT: Direct Microscopic Exam Of Tape Lift Samples

PROJECT NUMBER:

Sacramento, CA 95841

20804001

LABORATORY ID NUMBER:

0804013

Hygiene Technologies International, Inc.

Received Date:

April 07, 2008

Attention: Wes Frey

Report Date:

April 11, 2008

4330 Auburn Blvd. Suite 1850

Customer Sample Number	Date of Analysis	Method	Sample Intact	Amorphous Debris	Miscellaneous Fungi/Pollen ¹	Fungi with hyphal and /or sporulating structures ²	Loose spores/ Other comments ²
-TL01JL	04/11/08	M102.1	Yes	Light dander, Very light fibers, Very light particulates	Trace	None	Trace Epicoccum (single spore observed)
-TL02JL	04/11/08	M102.1	Yes	Moderate dander, Moderate fibers, Moderate particulates, Very light insect parts	Trace	None	Trace Cladosporium, Trace Curvularia*, Trace Epicoccum, Trace Oidium*, *(single spore observed)
-TL03JL	04/11/08	M102.1	Yes	Light dander, Light particulates, Very light fibers	Trace	None	None
-TL04JL	04/11/08	M102.1	Yes	Light dander, Light fibers, Very light particulates	Trace	None	None
-TL05JL	04/11/08	M102.1	Yes	Light dander, Light particulates, Very light fibers	Trace	None	Trace Cladosporium
-TL06JL	04/11/08	M102.1	Yes	Light dander, Very light fibers, Very light particulates	Trace	None	None
-TL07JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	None	None	None
-TL08JL	04/11/08	M102.1	Yes	Light dander, Very light fibers, Very light particulates	Trace	None	None
-TL09JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	Trace	None	Trace Penicillium/Aspergillus type:

^{1 -} Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

APPROVED:	DATE: 04/11/08
Name Lucas Wallin	Title: Lab Analyst

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.

^{2 -} Quantities of fungi are graded (from least to greatest) as a percentage of coverage of the slide area examined: none (0%), trace (0 - 10%), few (10 - 40%), numerous (40 - 80%), and massive (>80%).



2200 Amapola Ct., Suite 102 Torrance, CA 90501

(310) 618-0400 (310) 618-9400 fax

Page 2 of 3

FINAL REPORT: Direct Microscopic Exam Of Tape Lift Samples

PROJECT NUMBER: 20804001

LABORATORY ID NUMBER: 0804013

Hygiene Technologies International, Inc.

April 07, 2008

Attention: Wes Frey

Received Date: Report Date: April 11, 2008

4330 Auburn Blvd. Suite 1850

Sacramento, CA 95841

Customer Sample Number	Date of Analysis	Method	Sample Intact	Amorphous Debris	Miscellaneous Fungi/Pollen ¹	Fungi with hyphal and /or sporulating structures ²	Loose spores/ Other comments ²
-TL10JL	04/11/08	M102.1	Yes	Light dander, Light fibers, Light particulates	Trace	None	Trace Penicillium/Aspergillus types
-TL11JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	Trace	None	Trace Penicillium/Aspergillus types (single spore observed)
-TL12JL	04/11/08	M102.1	Yes	Moderate dander, Light fibers, Very light particulates	Trace	None	Trace Cladosporium, Trace Penicillium/Aspergillus types
-TL13JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	Trace	None	Trace Penicillium/Aspergillus types (single spore observed)
-TL14JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	None	None	None
-TL15JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	Trace	None	Trace Cladosporium (single spore observed)
-TL16JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	Trace	None	None
-TL17JL	04/11/08	M102.1	Yes	Very light dander, Very light fibers, Very light particulates	None	None	Trace Cladosporium (single spore observed)
-TL18JL	04/11/08	M102.1	Yes	Light dander, Light particulates, Very light fibers	Trace	None	Trace Penicillium/Aspergillus types

^{1 -} Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

APPROVED:		DATE:	d1/11/08
Name Lagas William	Title:	Lab Andyst	

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.

^{2 -} Quantities of fungi are graded (from least to greatest) as a percentage of coverage of the slide area examined: none (0%), trace (0 - 10%), few (10 - 40%), numerous (40 - 80%), and massive (>80%).



2200 Amapola Ct., Suite 102 Torrance, CA 90501

(310) 618-0400 (310) 618-9400 fax

Page 3 of 3

FINAL REPORT: Direct Microscopic Exam Of Tape Lift Samples

PROJECT NUMBER:

20804001

LABORATORY ID NUMBER:

0804013

Hygiene Technologies International, Inc.

Received Date:

April 07, 2008

Attention: Wes Frey

4330 Auburn Blvd. Suite 1850

Sacramento, CA 95841

Report Date: April 11, 2008

Customer Sample Number	Date of Analysis	Method	Sample Intact	Amorphous Debris	Miscellaneous Fungi/Pollen ¹	Fungi with hyphal and /or sporulating structures ²	Loose spores/ Other comments ²
-TL19JL	04/11/08	M102.1	Yes	Light dander, Light particulates, Very light fibers	Trace	None	Trace Cladosporium, Trace Penicillium/Aspergillus types, Trace Unidentified mitosporic fungi
-TL20JL	04/11/08	M102.1	Yes	Moderate dander, Moderate fibers, Light particulates	Trace	None	Trace Bipolaris/Drechslera group*, Trace Cladosporium, Trace Penicillium/Aspergillus types, Trace Torula*, Trace Ulocladium*, *(single spore observed)

APPROVED	: _/	200		DATE:	64/11/08	
Name	Lucas	Wallin	Title:	Lab Amlyst		

Results reported relate only to the sample items tested. This test report shall not be reproduced (except in full), corrected or added to without written approval from BioHygiene Laboratories, Inc.

^{1 -} Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating (indicative of normal trapping).

^{2 -} Quantities of fungi are graded (from least to greatest) as a percentage of coverage of the slide area examined: none (0%), trace (0 - 10%), few (10 - 40%), numerous (40 - 80%), and massive (>80%).